



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 45 /00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/03746</b> <b>(43) International Publication Date:</b> 27 January 2000 (27.01.00)
<b>(21) International Application Number:</b> PCT/US99/15827 <b>(22) International Filing Date:</b> 14 July 1999 (14.07.99)  <b>(30) Priority Data:</b> 09/115,387 14 July 1998 (14.07.98) US 09/273,224 19 March 1999 (19.03.99) US  <b>(71) Applicant:</b> THE BRIGHAM AND WOMEN'S HOSPITAL, INC. [US/US]; 75 Francis Street, Boston, MA 02115 (US). <b>(72) Inventor:</b> LIAO, James, K.; 12 Audubon Road, Weston, MA 02193 (US). <b>(74) Agent:</b> GATES, Edward, R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> UPREGULATION OF TYPE III ENDOTHELIAL CELL NITRIC OXIDE SYNTHASE BY AGENTS THAT DISRUPT ACTIN CYTOSKELETAL ORGANIZATION  <b>(57) Abstract</b> <p>A use for agents that disrupt actin cytoskeletal organization is provided. In the instant invention, agents that disrupt actin cytoskeletal organization are found to upregulate endothelial cell Nitric Oxide Synthase activity. As a result, agents that disrupt actin cytoskeletal organization are useful in treating or preventing conditions that result from the abnormally low expression and/or activity of endothelial cell Nitric Oxide Synthase. Such conditions include pulmonary hypertension, ischemic stroke, impotence, heart failure, hypoxia-induced conditions, insulin deficiency, progressive renal disease, gastric or esophageal motility syndrome, etc. Subjects thought to benefit mostly from such treatments include nonhyperlipidemics and nonhypercholesterolemics, but not necessarily exclude hyperlipidemics and hypercholesterolemics.</p> <div data-bbox="836 1165 1380 1512"><p>Western blot analysis showing protein levels. The top panel shows eNOS protein levels with bands at 205 kD and 117 kD. The bottom panel shows c-myc-RhoA protein levels with bands at 25 kD. Lanes are labeled c, wt, and N19.</p></div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

**UPREGULATION OF TYPE III ENDOTHELIAL CELL NITRIC OXIDE  
SYNTHASE BY  
AGENTS THAT DISRUPT ACTIN CYTOSKELETAL ORGANIZATION**

5

**Field of the Invention**

This invention relates to the use of agents that disrupt actin cytoskeletal organization as upregulators of Type III endothelial cell Nitric Oxide Synthase. Further, this invention relates to methods that employ agents that disrupt actin cytoskeletal organization to treat conditions that result from the abnormally low expression and/or activity of endothelial cell Nitric Oxide  
10 Synthase in a subject.

**Background of the Invention**

Nitric oxide (NO) has been recognized as an unusual messenger molecule with many physiologic roles, in the cardiovascular, neurologic and immune systems (Griffith, TM et al., *J*  
15 *Am Coll Cardiol*, 1988, 12:797-806). It mediates blood vessel relaxation, neurotransmission and pathogen suppression. NO is produced from the guanidino nitrogen of L-arginine by NO Synthase (Moncada, S and Higgs, EA. *Eur J Clin Invest*, 1991, 21(4):361-374). In mammals, at least three isoenzymes of NO Synthase have been identified. Two, expressed in neurons (nNOS) and endothelial cells (Type III-ecNOS), are calcium-dependent, whereas the third is calcium-  
20 independent and is expressed by macrophages and other cells after induction with cytokines (Type II-iNOS) (Bredt, DS and Snyder, SH, *Proc Natl Acad Sci USA*, 1990, 87:682-685, Janssens, SP et al., *J Biol Chem*, 1992, 267:22964, Lyons, CR et al., *J Biol Chem*, 1992, 267:6370-6374). The various physiological and pathological effects of NO can be explained by its reactivity and different routes of formation and metabolism.

25

Recent studies suggest that a loss of endothelial-derived NO activity may contribute to the atherogenic process (O'Driscoll, G, et al., *Circulation*, 1997, 95:1126-1131). For example, endothelial-derived NO inhibits several components of the atherogenic process including monocyte adhesion to the endothelial surface (Tsao, PS et al., *Circulation*, 1994, 89:2176-2182), platelet aggregation (Radomski, MW, et al., *Proc Natl Acad Sci USA*, 1990, 87:5193-5197),  
30 vascular smooth muscle cell proliferation (Garg, UC and Hassid, A, *J Clin Invest*, 1989, 83:1774-1777), and vasoconstriction (Tanner, FC et al., *Circulation*, 1991, 83:2012-2020). In addition, NO can prevent oxidative modification of low-density lipoprotein (LDL) which is a major contributor to atherosclerosis, particularly in its oxidized form (Cox, DA and Cohen, ML, *Pharm*

Rev, 1996, 48:3-19).

It has been shown in the prior art that hypoxia downregulates ecNOS expression and/or activity via decreases in both ecNOS gene transcription and mRNA stability (Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666, Shaul, PW et al., *Am J Physiol*, 1997, 272: L1005-L1012).  
5 Thus, ischemia-induced hypoxia may produce deleterious effects, in part, through decreases in ecNOS activity.

HMG-CoA (3-hydroxy-3-methylglutaryl-coenzyme A) reductase is the microsomal enzyme that catalyzes the rate limiting reaction in cholesterol biosynthesis (HMG-CoA6Mevalonate). An HMG-CoA reductase inhibitor inhibits HMG-CoA reductase, and  
10 as a result inhibits the synthesis of cholesterol. A number of HMG-CoA reductase inhibitors has been used to treat individuals with hypercholesterolemia. Clinical trials with such compounds have shown great reductions of cholesterol levels in hypercholesterolemic patients. Moreover, it has been shown that a reduction in serum cholesterol levels is correlated with improved endothelium-dependent relaxations in atherosclerotic vessels (Treasure, CB et al., *N Engl J Med*,  
15 1995, 332:481-487). Indeed, one of the earliest recognizable benefits after treatment with HMG-CoA reductase inhibitors is the restoration of endothelium-dependent relaxations (*supra*, Anderson, TJ et al., *N Engl J Med*, 1995, 332:488-493).

Although the mechanism by which HMG-CoA reductase inhibitors restore endothelial function is primarily attributed to the inhibition of hepatic HMG-CoA reductase and the  
20 subsequent lowering of serum cholesterol levels, little is known on whether inhibition of endothelial HMG-CoA reductase has additional beneficial effects on endothelial function.

By inhibiting L-mevalonate synthesis, HMG-CoA reductase inhibitors also prevent the synthesis of other important isoprenoid intermediates of the cholesterol biosynthetic pathway, such as farnesylpyrophosphate (FPP) and geranylgeranylpyrophosphate (GGPP) (Goldstein, JL  
25 and Brown, MS, *Nature*, 1990, 343:425-430). The isoprenoids are important lipid attachments for the post-translational modification of variety of proteins, including G-protein and G-protein subunits, Heme-a, nuclear lamins, and Ras-like proteins, such as Rho, Rab, Rac, Ral or Rap (Goldstein, JL and Brown, MS, *supra*; Casey, PJ, *Science*, 1995, 268:221-225). The role that isoprenoids play in regulating ecNOS expression, however, is not known.

30 Pulmonary hypertension is a major cause of morbidity and mortality in individuals exposed to hypoxic conditions (Scherrer, U et al., *N Engl J Med*, 1996, 334:624-629). Recent studies demonstrate that pulmonary arterial vessels from patients with pulmonary hypertension

-3-

have impaired release of NO (Giaid, A and Saleh, D, *N Engl J Med*, 1995, 333:214-221, Shaul, PW, *Am J Physiol*, 1997, 272: L1005-L1012). Additionally, individuals with pulmonary hypertension demonstrate reduced levels of ecNOS expression in their pulmonary vessels and benefit clinically from inhalation nitric oxide therapy (Roberts, JD et al., *N Engl J Med*, 1997, 336:605-610, Kouyoumdjian, C et al., *J Clin Invest*, 1994, 94:578-584). Conversely, mutant mice lacking ecNOS gene or newborn lambs treated with the ecNOS inhibitor, N-omega-monomethyl-L-arginine (LNMA, or N-omega-nitro-L-arginine), develop progressive elevation of pulmonary arterial pressures and resistance (Steudel, W et al., *Circ Res*, 1997, 81:34-41, Fineman, JR et al., *J Clin Invest*, 1994, 93:2675-2683). It has also been shown in the prior art that hypoxia causes pulmonary vasoconstriction via inhibition of endothelial cell nitric oxide synthase (ecNOS) expression and activity (Adnot, S et al., *J Clin Invest*, 1991, 87:155-162, Liao, JK et al., *J Clin Invest*, 1995, 96, 2661-2666). Hence, hypoxia-mediated downregulation of ecNOS may lead to the vasoconstrictive and structural changes associated with pulmonary hypertension.

Often cited as the third most frequent cause of death in the developed countries, stroke has been defined as the abrupt impairment of brain function caused by a variety of pathologic changes involving one or several intracranial or extracranial blood vessels. Approximately 80% of all strokes are ischemic strokes, resulting from restricted blood flow. Mutant mice lacking the gene for ecNOS are hypertensive (Huang, PL et al., *Nature*, 1995, 377:239-242, Steudel, W et al., *Circ Res*, 1997, 81:34-41) and develop greater intimal smooth muscle proliferation in response to cuff injury. Furthermore, occlusion of the middle cerebral artery results in 21% greater infarct size in "ecNOS knockout" mice compared to wildtype mice (Huang, Z et al., *J Cereb Blood Flow Metab*, 1996, 16:981-987). These findings suggest that the ecNOS production may play a role in cerebral infarct formation and sizes. Additionally, since most patients with ischemic strokes have average or normal cholesterol levels, little is known on what the potential benefits of HMG-CoA reductase inhibitor administration would be in cerebrovascular events.

There exists a need to identify agents that improve endothelial cell function.

There also exists a need to identify agents that can be used acutely or in a prophylactic manner to treat conditions that result from low levels of endothelial cell Nitric Oxide Synthase.

### Summary of the Invention

The invention involves the discovery that agents which disrupt actin cytoskeletal

-4-

organization can upregulate endothelial cell Nitric Oxide Synthase (Type III) expression. The invention, therefore, is useful whenever it is desirable to restore endothelial cell Nitric Oxide Synthase activity or increase such activity in a cell, tissue or subject, provided the cell or the tissue expresses endothelial cell Nitric Oxide Synthase.

5 Nitric Oxide Synthase activity is involved in many conditions, including impotence, heart failure, gastric and esophageal motility disorders, kidney disorders such as kidney hypertension and progressive renal disease, insulin deficiency, etc. Individuals with such conditions would benefit from increased endothelial cell Nitric Oxide Synthase activity. It also was known that individuals with pulmonary hypertension demonstrate reduced levels of Nitric Oxide Synthase  
10 expression in their pulmonary vessels and benefit clinically from inhalation of Nitric Oxide. The invention therefore is particularly useful for treating pulmonary hypertension. It also has been demonstrated that hypoxia causes an inhibition of endothelial cell Nitric Oxide Synthase activity. The invention therefore is useful for treating subjects with hypoxia-induced conditions. It also has been discovered, surprisingly, that agents which disrupt actin cytoskeletal organization are  
15 useful for reducing brain injury that occurs following a stroke.

According to one aspect of the invention, a method is provided for increasing endothelial cell Nitric Oxide Synthase activity in a subject who would benefit from increased endothelial cell Nitric Oxide Synthase activity in a tissue. The method involves administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization that increases endothelial  
20 cell Nitric Oxide Synthase activity in an amount(s) effective to increase endothelial cell Nitric Oxide Synthase activity in the tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

In certain embodiments, the agent that disrupt actin cytoskeletal organization is not H-7 [1-(5-isoquinoline sulphonyl)-2-methylpiperazine dihydro-chloride], when the subject in need  
25 of such treatment has an abnormally elevated risk of an ischemic stroke because of having experienced a previous ischemic event, or has experienced an ischemic stroke. In other embodiments, the agent that disrupt actin cytoskeletal organization is not Fasudil (HA1077) [Hexahydro-1-(5-isoquinolinesulphonyl)-1H-1,4- diazepine] when the subject in need of such treatment has an abnormally elevated risk of stroke, or has experienced stroke.

30 In one important embodiment, agents that disrupt actin cytoskeletal organization do not affect cholesterol levels in a subject. In certain embodiments, however, agents that disrupt actin cytoskeletal organization as well as increasing endothelial cell Nitric Oxide Synthase activity in

-5-

the tissue of a subject can also affect cholesterol levels in the subject. In atherosclerotic patients, reduction in serum cholesterol is correlated with improved endothelium-dependent relaxation in atherosclerotic vessels (Treasure, et al., *N.Engl.J.Med.*, 1995, 332:481-487). HMG-CoA reductase inhibitors have been demonstrated to reduce serum cholesterol in a matter of weeks, and maximum level of cholesterol reduction can be achieved after a few months of chronic administration. In contrast, the effect of agents that disrupt actin cytoskeletal organization on up-regulation of eNOS occurs within a few days. Thus, treatment according to the present invention provides significant advantages, e.g., when administered to address short term increases in risk of stroke or other embolic events, such as that due to surgical intervention, even for hypercholesterolemic patients.

In certain embodiments, the subject is not hypercholesterolemic or not hypertriglyceridemic or both (i.e., nonhyperlipidemic). In other embodiments the amount is sufficient to increase endothelial cell Nitric Oxide Synthase activity above normal baseline levels established by age-controlled groups, described in greater detail below.

The subject can have a condition characterized by an abnormally low level of endothelial cell Nitric Oxide Synthase activity which is hypoxia-induced. In other embodiments the subject can have a condition comprising an abnormally low level of endothelial cell Nitric Oxide Synthase activity which is chemically induced. In still other embodiments the subject can have a condition comprising an abnormally low level of endothelial cell Nitric Oxide Synthase activity which is cytokine induced. In certain important embodiments, the subject has pulmonary hypertension or an abnormally elevated risk of pulmonary hypertension. In other important embodiments, the subject has experienced an ischemic stroke or has an abnormally elevated risk of an ischemic stroke. In still other important embodiments, the subject has heart failure or progressive renal disease. In yet other important embodiments, the subject is chronically exposed to hypoxic conditions.

In further important embodiments, the subject has experienced a thrombotic event or has an abnormally elevated risk of thrombosis. In still other embodiments, the subject has an abnormally elevated risk of arteriosclerosis or has arteriosclerosis. In other important embodiments, the subject has an abnormally elevated risk of developing a myocardial infarction or has experienced a myocardial infarction. In yet further important embodiments, the subject has an abnormally elevated risk of reperfusion injury. In preferred embodiments, the subject with an elevated risk of reperfusion injury is an organ transplant recipient (e.g., heart, kidney, liver,

-6-

etc.). In other important embodiments, the subject has homocystinuria. In certain other important embodiments, the subject has Cerebral autosomal dominant arteriopathy with subcortical infarcts and leucoencephalopathy (CADASIL) syndrome. In further important embodiments, the subject has a degenerative disorder of the nervous system. In preferred embodiments, the subject with  
5 a degenerative disorder of the nervous system has Alzheimer's disease.

According to any of the foregoing embodiments, the preferred agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D. In some embodiments the myosin light chain  
10 kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene- 1-sulphonyl)-1H- hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl) -2-methylpiperazine dihydro-chloride. Likewise, in any of the foregoing embodiments, the method can further comprise co-administering an endothelial cell Nitric Oxide Synthase substrate (L-arginine preferred) and/or co-administering an agent other  
15 than an agent that disrupts actin cytoskeletal organization that also increases endothelial cell Nitric Oxide Synthase activity, and/or co-administering at least one different agent that disrupts actin cytoskeletal organization. A preferred agent other than an agent that disrupts actin cytoskeletal organization is selected from the group consisting of estrogens and angiotensin-converting enzyme (ACE) inhibitors. The agents may be administered to a subject who has a  
20 condition or prophylactically to a subject who has a risk, and more preferably, an abnormally elevated risk, of developing a condition. The inhibitors also may be administered acutely.

According to another aspect of the invention, a method is provided for increasing endothelial cell Nitric Oxide Synthase activity in a subject to treat a condition favorably affected by an increase in endothelial cell Nitric Oxide Synthase activity in a tissue. Such conditions are  
25 exemplified above. The method involves administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor. In important embodiments, agents that disrupt actin cytoskeletal organization do not affect cholesterol levels  
30 in a subject. In certain embodiments, however, agents that disrupt actin cytoskeletal organization as well as increase endothelial cell Nitric Oxide Synthase activity in the tissue of a subject can also affect cholesterol levels in the subject. In certain embodiments, the subject is



-7-

nonhypercholesterolemic. In other embodiments, the subject is nonhyperlipidemic. Important conditions are as described above. Also as described above, the method can involve co-administration of substrates of endothelial cell Nitric Oxide Synthase and/or co-administering an agent other than an agent that disrupts actin cytoskeletal organization that also increases  
5 endothelial cell Nitric Oxide Synthase activity, and/or co-administering at least one different agent that disrupts actin cytoskeletal organization. Preferred compounds are as described above. As above, the agents that disrupt actin cytoskeletal organization with or without the co-administered compounds can be administered, *inter alia*, acutely or prophylactically.

According to another aspect of the invention, a method is provided for reducing brain  
10 injury resulting from stroke. The method involves administering to a subject having an abnormally high risk of an ischemic stroke an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the brain of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor. As above, important embodiments include the agent being selected  
15 from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D. As above, in some embodiments a myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-1H- hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)  
20 -2-methylpiperazine dihydro-chloride. Also as above, important embodiments include co-administering a substrate of endothelial cell Nitric Oxide Synthase (L-arginine preferred), and/or co-administering an agent other than an agent that disrupts actin cytoskeletal organization that also increases endothelial cell Nitric Oxide Synthase activity, and/or co-administering at least one different agent that disrupts actin cytoskeletal organization. Likewise, important embodiments  
25 include prophylactic and acute administration of the agent(s).

According to another aspect of the invention, a method is provided for treating pulmonary hypertension. The method involves administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase pulmonary endothelial cell Nitric Oxide Synthase activity in the subject, provided that the agent that disrupts  
30 actin cytoskeletal organization is not a *rho* GTPase function inhibitor. Particularly important embodiments are as described above in connection with the methods for treating brain injury. Another important embodiment is administering the agent prophylactically to a subject who has

-8-

an abnormally elevated risk of developing pulmonary hypertension, including subjects that are chronically exposed to hypoxic conditions.

According to another aspect of the invention, a method for treating heart failure is provided. The method involves administering to a subject in need of such treatment an agent that  
5 disrupts actin cytoskeletal organization in an amount effective to increase vascular endothelial cell Nitric Oxide Synthase activity in the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor. As discussed above, important embodiments include prophylactic and acute administration of the agent(s). Preferred compounds and co-administration schemes are as described above.

10 According to yet another aspect of the invention, a method is provided for treating progressive renal disease. The method involves administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase renal endothelial cell Nitric Oxide Synthase activity in the kidney of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.  
15 Important embodiments and preferred compounds and schemes of co-administration are as described above in connection with heart failure.

According to another aspect of the invention, a method for increasing blood flow in a tissue of a subject is provided. The method involves administering to a subject in need of such treatment a first agent that disrupts actin cytoskeletal organization in an amount effective to  
20 increase endothelial cell Nitric Oxide Synthase activity in the tissue of the subject, provided that the first agent is not an agent selected from the group consisting of a *rho* GTPase function inhibitor, H7, and fasudil. In preferred embodiments, the tissue in which blood flow is increased includes tissue in the brain. In a particularly preferred embodiment, cerebral blood flow is enhanced. In certain embodiments the first agent is not a myosin light chain kinase inhibitor.  
25 In other embodiments the first agent is selected from the group consisting of a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D. Other important embodiments include co-administering a second agent to the subject with a condition treatable by the second agent in an amount effective to treat the condition, whereby the delivery of the second agent to a tissue of the subject is enhanced as a  
30 result of the increased blood flow. In certain embodiments, the tissue is brain and the second agent comprises an agent having a site of action in the brain.

According to another aspect of the invention, a method of screening for identifying an

-9-

agent that disrupts actin cytoskeletal organization for treatment of subjects who would benefit from increased endothelial cell Nitric Oxide Synthase activity in a tissue, is provided. The method involves identifying an agent that disrupts actin cytoskeletal organization suspected of increasing endothelial cell Nitric Oxide Synthase activity, and determining whether or not the agent that disrupts actin cytoskeletal organization produces an increase in endothelial cell Nitric Oxide Synthase activity *in vivo* or *in vitro*. In certain embodiments, the subject who would benefit from increased endothelial cell Nitric Oxide Synthase activity has an abnormally elevated risk of stroke.

The invention also involves the use of agents that disrupt actin cytoskeletal organization in the manufacture of medicaments for treating the above-noted conditions. Important conditions, compounds, etc. are as described above. The invention further involves pharmaceutical preparations including the agents that disrupt actin cytoskeletal organization for treating the above-noted conditions. The preparations can include other agents such as second agents, ecNOS substrates, ecNOS cofactors, as described above, or can be cocktails of agents that disrupt actin cytoskeletal organization according to the invention [non-*rho* GTPase function inhibitor(s)]. In certain embodiments, however, the cocktails can include a *rho* GTPase function inhibitor(s) that disrupts actin cytoskeletal organization together with the agents that disrupt actin cytoskeletal organization agent of the invention. The invention also involves pharmaceutical preparations that are cocktails of agents that disrupt actin cytoskeletal organization together with agents other than agents that disrupt actin cytoskeletal organization that also increase ecNOS activity in a cell, directly or indirectly (synergistically, cooperatively, additively, etc.).

In certain embodiments, compositions and pharmaceutical preparations that are cocktails of an agent that disrupt actin cytoskeletal organization and L-arginine are provided. In other embodiments, the agent that disrupt actin cytoskeletal organization and the L-arginine are in amounts effective to increase blood flow. In further embodiments, the agent that disrupt actin cytoskeletal organization and the L-arginine are in amounts effective to increase blood flow in brain tissue. In preferred embodiments, administration of the agent that disrupt actin cytoskeletal organization and the L-arginine results in increased blood flow. In particularly preferred embodiments, administration of the agent that disrupt actin cytoskeletal organization and the L-arginine results in increased blood flow to the brain. Any of the above cocktail compositions may also include other cofactors that enhance ecNOS substrate conversion by ecNOS to nitric oxide, the preferred cofactors being NADPH and tetrahydrobiopterin.

-10-

The invention also involves methods for increasing ecNOS activity in a cell by contacting the cell with an effective amount of a agent that disrupt actin cytoskeletal organization (excluding *rho* GTPase function inhibitors), alone, or together with any of the agents co-administered as described above, or as a cocktail as described above. Any of the above cocktails may also include  
5 a substrate for endothelial cell Nitric Oxide Synthase, the preferred substrate being L-arginine, and/or other cofactors that enhance ecNOS substrate conversion by ecNOS to nitric oxide, the preferred cofactors being NADPH and tetrahydrobiopterin.

In any of the foregoing aspects of the invention the agent can be a non-fasudil agent that disrupts actin cytoskeletal organization.

10 These and other aspects of the invention are described in greater detail below.

#### **Brief Description of the Drawings**

Figure 1. ecNOS activity and expression in wild-type SV-129 mice aortas with and without treatment with simvastatin for 14 days.

15 Figure 2. ecNOS mRNA expression in the infarcted, ipsilateral (I) and not-infarcted, contralateral (C) forebrain hemispheres of SV-129 mice with and without treatment with simvastatin.

Figure 3. Northern blots showing the effects of mevastatin alone or in combination with FPP or GGPP on eNOS (ecNOS) steady-state mRNA levels after 24h.

20 Figure 4. Western blots showing the effects of mevastatin alone or in combination with FPP or GGPP or LDL-cholesterol on eNOS (ecNOS) protein levels after 24h.

Figure 5. Western blots showing the effects of C3 transferase, mevastatin, or L-mevalonate on eNOS (ecNOS) protein levels after 24h.

Figure 6. Western blots showing eNOS (ecNOS) protein levels after transfection with  
25 insertless vector, pcDNA3 (C), c-myc-wildtype-RhoA (wt), and c-myc-N19RhoA (dominant-negative rhoA mutant).

Figure 7. Effects of C3 transferase, FPP, GGPP, and CNF-1 on mevastatin-induced eNOS (ecNOS) activity as determined by LNMA-inhibitable nitrite production at 24 h.

Figure 8. Immunoblots showing the concentration-dependent effects of MLC kinase  
30 inhibitor H-7on ecNOS protein levels after 24 hours.

Figure 9. Northern blots showing ecNOS expression of endothelial cells treated with cytochalasin D at 24 hours.

-11-

Figure 10. Immunoblots showing the concentration-dependent effects of 2, 3-butanedione 2-monoxime on ecNOS protein levels.

Figure 11. Northern blots showing ecNOS expression of endothelial cells treated with nocodazole for 24 hours.

5        Figure 12. Bar graph showing regional CBF changes in wild type and eNOS null mice for 40 min after L-arginine or saline infusion.

Figure 13. Bar graph showing regional CBF changes in simvastatin-treated mice for 40 min after L-arginine or saline infusion at the same dose.

10

### **Detailed Description of the Invention**

The invention is useful whenever it is desirable to increase endothelial cell Nitric Oxide Synthase (Type III isoform) activity in a cell, in a tissue, or in a subject. A subject as used herein includes humans, non human primates, dogs, cats, sheep, goats, cows, pigs, horses and rodents. The invention thus is useful for therapeutic purposes and also is useful for research purposes such as in testing in animal or *in vitro* models of medical, physiological or metabolic pathways or conditions. Nitric Oxide Synthase is the enzyme that catalyzes the reaction that produces nitric oxide from the substrate L-arginine. As the name implies, endothelial cell nitric oxide Synthase refers to the Type III isoform of the enzyme found in the endothelium.

By "ecNOS activity", it is meant the ability of a cell to generate nitric oxide from the substrate L-arginine. Increased ecNOS activity can be accomplished in a number of different ways. For example, an increase in the amount of ecNOS protein or an increase in the activity of the protein (while maintaining a constant level of the protein) can result in increased "activity". An increase in the amount of protein available can result from increased transcription of the ecNOS gene, increased stability of the ecNOS mRNA or a decrease in ecNOS protein degradation. (The term "expression" is used interchangeably with the term "activity" throughout this application).

The ecNOS activity in a cell or in a tissue can be measured in a variety of different ways. A direct measure would be to measure the amount of ecNOS present. Another direct measure would be to measure the amount of conversion of arginine to citrulline by ecNOS or the amount of generation of nitric oxide by ecNOS under particular conditions, such as the physiologic conditions of the tissue. The ecNOS activity also can be measured more indirectly, for example by measuring mRNA half-life (an upstream indicator) or by a phenotypic response to the

-12-

presence of nitric oxide (a downstream indicator). One phenotypic measurement employed in the art is detecting endothelial dependent relaxation in response to a acetylcholine, which response is affected by ecNOS activity. The level of nitric oxide present in a sample can be measured using a nitric oxide meter. All of the foregoing techniques are well known to those of ordinary skill in the art, and some are described in the examples below.

The present invention, by causing an increase in ecNOS activity, permits not only the re-establishment of normal base-line levels of ecNOS activity, but also allows increasing such activity above normal base-line levels. Normal base-line levels are the amounts of activity in a normal control group, controlled for age and having no symptoms which would indicate alteration of endothelial cell Nitric Oxide Synthase activity (such as hypoxic conditions, hyperlipidemia and the like). The actual level then will depend upon the particular age group selected and the particular measure employed to assay activity. Specific examples of various measures are provided below. In abnormal circumstances, e.g. hypoxic conditions, pulmonary hypertension, etc., endothelial cell Nitric Oxide Synthase activity is depressed below normal levels. Surprisingly, when using agents that disrupt actin cytoskeletal organization according to the invention, not only can normal base-line levels be restored in such abnormal conditions, but endothelial cell Nitric Oxide Synthase activity can be increased desirably far above normal base-line levels of endothelial cell Nitric Oxide Synthase activity. Thus, "increasing activity" means any increase in endothelial cell Nitric Oxide Synthase activity in the subject resulting from the treatment with agents that disrupt actin cytoskeletal organization according to the invention, including, but not limited to, such activity as would be sufficient to restore normal base-line levels and such activity as would be sufficient to elevate the activity above normal base-line levels.

As mentioned above, Nitric Oxide Synthase activity is involved in many conditions, including stroke, pulmonary hypertension, thrombosis, arteriosclerosis, myocardial infarction, reperfusion injury (e.g., in an organ transplant recipient), impotence, heart failure, gastric and esophageal motility disorders, kidney disorders such as kidney hypertension and progressive renal disease, insulin deficiency, hypoxia-induced conditions, homocystinuria, neurodegenerative disorders, CADASIL syndrome, etc. In one embodiment of the invention the decrease in endothelial cell Nitric Oxide Synthase activity is cytokine induced. Cytokines are soluble polypeptides produced by a wide variety of cells that control gene activation and cell surface molecule expression. They play an essential role in the development of the immune system and

-13-

thus in the development of an immune response. However, besides their numerous beneficial properties, they have also been implicated in the mechanisms for the development of a variety of inflammatory diseases. For example, the cytokines TNF- $\alpha$  and IL-1 are thought to be part of the disease causing mechanism of non-cholesterol induced atherosclerosis, transplant arterial  
5 sclerosis, rheumatoid arthritis, lupus, scleroderma, emphysema, etc. Subjects of such disorders exhibit lower levels of endothelial cell Nitric Oxide Synthase activity (which is thus "cytokine induced"), and may benefit from therapy using the agents of the instant invention.

One important embodiment of the invention is treatment of ischemic stroke. Ischemic stroke (ischemic cerebral infarction) is an acute neurologic injury that results from a decrease in  
10 the blood flow involving the blood vessels of the brain. Ischemic stroke is divided into two broad categories, thrombotic and embolic.

A surprising finding was made in connection with the treatment of ischemic stroke. In particular, it was discovered that treatment according to the invention can reduce the brain injury that follows an ischemic stroke. Brain injury reduction, as demonstrated in the examples below,  
15 can be measured by determining a reduction in infarct size in the treated versus the control groups. Likewise, functional tests measuring neurological deficits provided further evidence of reduction in brain injury in the treated animals versus the controls. Cerebral blood flow also was better in the treated animals versus the controls. Thus, in the various accepted models of brain injury following stroke, a positive effect was observed in the treated animals versus the control  
20 animals. It is believed that all of the foregoing positive results are attributable to the upregulation of endothelial cell Nitric Oxide Synthase activity, which is believed demonstrated in the examples below.

An important embodiment of the invention is treatment of a subject with an abnormally elevated risk of an ischemic stroke. As used herein, subjects having an abnormally elevated risk  
25 of an ischemic stroke are a category determined according to conventional medical practice; such subjects may also be identified in conventional medical practice as having known risk factors for stroke or having increased risk of cerebrovascular events. This category includes, for example, subjects which are having elected vascular surgery. Typically, the risk factors associated with cardiac disease are the same as are associated with stroke. The primary risk factors include  
30 hypertension, hypercholesterolemia, and smoking. In addition, atrial fibrillation or recent myocardial infarction are important risk factors.

As used herein, subjects having an abnormally elevated risk of an ischemic stroke also

-14-

include individuals undergoing surgical or diagnostic procedures which risk release of emboli, lowering of blood pressure or decrease in blood flow to the brain, such as carotid endarterectomy, brain angiography, neurosurgical procedures in which blood vessels are compressed or occluded, cardiac catheterization, angioplasty, including balloon angioplasty, coronary by-pass surgery, or similar procedures. Subjects having an abnormally elevated risk of an ischemic stroke also include individuals having any cardiac condition that may lead to decreased blood flow to the brain, such as atrial fibrillation, ventricular tachycardia, dilated cardiomyopathy and other cardiac conditions requiring anticoagulation. Subjects having an abnormally elevated risk of an ischemic stroke also include individuals having conditions including arteriopathy or brain vasculitis, such as that caused by lupus, congenital diseases of blood vessels, such as cadasil syndrome, or migraine, especially prolonged episodes. In certain embodiments, the subject is not hypercholesterolemic or not hypertriglyceridemic or both (i.e., nonhyperlipidemic).

The treatment of stroke can be for patients who have experienced a stroke or can be a prophylactic treatment. Short term prophylactic treatment is indicated for subjects having surgical or diagnostic procedures which risk release of emboli, lowering of blood pressure or decrease in blood flow to the brain, to reduce the injury due to any ischemic event that occurs as a consequence of the procedure. Longer term or chronic prophylactic treatment is indicated for subjects having cardiac conditions that may lead to decreased blood flow to the brain, or conditions directly affecting brain vasculature. If prophylactic, then the treatment is for subjects having an abnormally elevated risk of an ischemic stroke, as described above. If the subject has experienced a stroke, then the treatment can include acute treatment. Acute treatment for stroke subjects means administration of the agents that disrupt actin cytoskeletal organization at the onset of symptoms of the condition or at the onset of a substantial change in the symptoms of an existing condition.

Another important embodiment of the invention is treatment of pulmonary hypertension. Pulmonary hypertension is a disease characterized by increased pulmonary arterial pressure and pulmonary vascular resistance. Hypoxemia, hypocapnia, and an abnormal diffusing capacity for carbon monoxide are almost invariable findings of the disease. Additionally, according to the present invention, patients with pulmonary hypertension also have reduced levels of ecNOS expression and/or activity in their pulmonary vessels. Traditionally, the criteria for subjects with, or at risk for pulmonary hypertension are defined on the basis of clinical and histological characteristics according to Heath and Edwards (*Circulation*, 1958, 18:533-547).



-15-

Subjects may be treated prophylactically to reduce the risk of pulmonary hypertension or subjects with pulmonary hypertension may be treated long term and/or acutely. If the treatment is prophylactic, then the subjects treated are those with an abnormally elevated risk of pulmonary hypertension. A subject with an abnormally elevated risk of pulmonary hypertension is a subject  
5 with chronic exposure to hypoxic conditions, a subject with sustained vasoconstriction, a subject with multiple pulmonary emboli, a subject with cardiomegaly and/or a subject with a family history of pulmonary hypertension.

Another important embodiment of the invention involves treating hypoxia-induced conditions. Hypoxia as used herein is defined as the decrease below normal levels of oxygen in  
10 a tissue. Hypoxia can result from a variety of circumstances, but most frequently results from impaired lung function. Impaired lung function can be caused by emphysema, cigarette smoking, chronic bronchitis, asthma, infectious agents, pneumonitis (infectious or chemical), lupus, rheumatoid arthritis, inherited disorders such as cystic fibrosis, obesity,  $\alpha_1$ -antitrypsin deficiency and the like. It also can result from non-lung impairments such as from living at very high  
15 altitudes. Hypoxia can result in pulmonary vasoconstriction via inhibition of eNOS activity.

Another important embodiment of the invention is the treatment of heart failure. Heart failure is a clinical syndrome of diverse etiologies linked by the common denominator of impaired heart pumping and is characterized by the failure of the heart to pump blood commensurate with the requirements of the metabolizing tissues, or to do so only from an  
20 elevating filling pressure.

In certain aspects of the invention, agents that disrupt actin cytoskeletal organization are administered to subjects that would benefit from increased endothelial cell Nitric Oxide Synthase activity. The administration of one or more agents that disrupt actin cytoskeletal organization is in an amount(s) effective to increase endothelial cell Nitric Oxide Synthase activity in tissue of  
25 the subject, provided that the agent that disrupts actin cytoskeletal organization used is not a *rho* GTPase function inhibitor (See later discussion). In certain embodiments, the subject is both nonhypercholesterolemic and/or nonhypertriglyceridemic, i.e., nonhyperlipidemic. Such subjects are thought to benefit mostly from the treatments of the invention, but the treatments do not necessarily exclude hyperlipidemic and hypercholesterolemic subjects.

30 A nonhypercholesterolemic subject is one that does not fit the current criteria established for a hypercholesterolemic subject. A nonhypertriglyceridemic subject is one that does not fit the current criteria established for a hypertriglyceridemic subject (See, e.g., Harrison's Principles of

-16-

Experimental Medicine, 13th Edition, McGraw-Hill, Inc., N.Y.- hereinafter "Harrison's"). Hypercholesterolemic subjects and hypertriglyceridemic subjects are associated with increased incidence of premature coronary heart disease. A hypercholesterolemic subject has an LDL level of >160 mg/dL or >130 mg/dL and at least two risk factors selected from the group consisting of male gender, family history of premature coronary heart disease, cigarette smoking (more than 10 per day), hypertension, low HDL (<35 mg/dL), diabetes mellitus, hyperinsulinemia, abdominal obesity, high lipoprotein (a), and personal history of cerebrovascular disease or occlusive peripheral vascular disease. A hypertriglyceridemic subject has a triglyceride (TG) level of >250 mg/dL. Thus, a hyperlipidemic subject is defined as one whose cholesterol and triglyceride levels equal or exceed the limits set as described above for both the hypercholesterolemic and hypertriglyceridemic subjects.

Another important embodiment of the invention is treatment of thrombosis. Thromboembolism is the collective term used for diseases characterized by the formation, development, or presence of a thrombus and the blocking of a vessel by a thrombus brought to a thrombotic vascular site by the blood current. Thromboembolism can reduce blood flow to almost all organs including the brain and myocardium. Thromboembolism involving the brain is otherwise known as an ischemic stroke and is described elsewhere in this application. Thromboembolism involving the heart is otherwise known as a myocardial infarction and is also described elsewhere in this application. According to Harrison's, certain patient groups have been identified who are particularly prone to thrombosis and embolism. These include patients: (1) immobilized after surgery; (2) with chronic congestive heart failure; (3) with atherosclerotic vascular disease; (4) with malignancy; or (5) who are pregnant.

An important embodiment of the invention is treatment of subjects with an abnormally elevated risk of thrombosis (or thromboembolism). As used herein, subjects having an abnormally elevated risk of thrombosis are a category determined according to conventional medical practice. Typically, prethrombotic patients can be identified by a careful history. There are, according to Harrison's, three important clues to this diagnosis: (1) repeated episodes of thromboembolism without an obvious predisposing condition; (2) a family history of thrombosis; and (3) well-documented thromboembolism in adolescents and young adults. Subjects may be treated prophylactically to reduce the risk of a thrombotic episode or subjects with thrombosis may be treated long-term and/or acutely. If the treatment is prophylactic, then the subjects treated are those with an abnormally elevated risk of thrombosis.

Another important embodiment of the invention is treatment of myocardial infarction. Myocardial infarction is the diseased state which occurs with the abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by atherosclerosis. Such injury is produced or facilitated by factors such as cigarette smoking, hypertension and lipid accumulation.

An important embodiment of the invention is treatment of a subject with an abnormally elevated risk of myocardial infarction. As used herein, subjects having an abnormally elevated risk of myocardial infarction are the category of patients that include those with unstable angina, multiple coronary risk factors (similar to those described for stroke elsewhere herein), and Prinzmetal's variant angina. Less common etiologic factors include hypercoagulability, coronary emboli, collagen vascular disease, and cocaine abuse.

Subjects may be treated prophylactically to reduce the risk of myocardial infarction, or subjects with myocardial infarction, may be treated long-term and/or acutely. If the treatment is prophylactic, then the subjects treated are those with an abnormally elevated risk of myocardial infarction. A subject with an abnormally elevated risk of myocardial infarction is a subject that falls in the above-described categories.

Another important embodiment of the invention, is the treatment of subjects with an abnormally elevated risk of reperfusion injury damage. Preferred subjects are about to receive or have received a transplant. According to the present invention, increase in eNOS expression and/or activity in the vessels of the transplanted organ is believed to reduce reperfusion injury damage. Reperfusion injury is the functional, metabolic, or structural change that includes necrosis in ischemic tissues, thought to result from reperfusion to ischemic areas of the tissue. The most common example involves myocardial reperfusion injury. In myocardial reperfusion injury, changes in ischemic heart muscle are thought to result from reperfusion to the ischemic areas of the heart. Changes can be fatal to muscle cells and may include oedema with explosive cell swelling and disintegration, sarcolemma disruption, fragmentation of mitochondria, contraction and necrosis, enzyme washout and calcium overload.

Another important embodiment of the invention, is the treatment of subjects with a homocystinuria. The homocystinurias are seven biochemically and clinically distinct disorders, each characterized by increased concentration of the sulfur-containing amino acid homocysteine in blood and urine. This is because the enzyme cystathione synthetase that converts homocysteine

and serine into cystathione, a precursor of cysteine, is missing. Subjects with a homocystinuria are also likely to suffer from thrombosis, and can benefit from increased ecNOS expression and/or activity.

Another important embodiment of the invention, is the treatment of subjects with Cerebral autosomal dominant arteriopathy with subcortical infarcts and leucoencephalopathy (CADASIL) syndrome. The disorder is characterized by relapsing strokes with neuropsychiatric symptoms and affects relatively young adults of both sexes. CT scans have demonstrated occlusive cerebrovascular infarcts in the white matter, which was usually reduced. Subjects with CADASIL syndrome can also benefit from increased ecNOS expression and/or activity.

Another important embodiment of the invention, is the treatment of subjects with a neurodegenerative disease. The term "neurodegenerative disease" is meant to include any pathological state involving neuronal degeneration, including Parkinson's Disease, Huntington's Disease, Alzheimer's Disease, and amyotrophic lateral sclerosis (ALS). In preferred embodiments, the neurodegenerative disease is Alzheimer's Disease. Alzheimer's Disease is a progressive, neurodegenerative disease characterised by loss of function and death of nerve cells in several areas of the brain leading to loss of cognitive function such as memory and language. The cause of nerve cell death is unknown but the cells are recognised by the appearance of unusual helical protein filaments in the nerve cells (neurofibrillary tangles) and by degeneration in cortical regions of brain, especially frontal and temporal lobes. Increase of cerebral blood flow mediated by an increase in ecNOS expression and/or activity can also be of benefit to subjects suffering from a neurodegenerative disease.

The invention involves treatment of the foregoing conditions using agents that disrupt actin cytoskeletal organization. Actin comprises a large proportion of the cytoplasmic proteins of many cells. Actin is present primarily in its globular form (G-actin), a single polypeptide 375 amino acids long, and is associated with one molecule of non-covalently bound ATP. The terminal phosphate of the ATP is hydrolysed after the actin polymerizes to form actin filaments (fibrous actin or F-actin). Actin filaments consist of a tight-helix of uniformly oriented actin monomers. They are polar structures, with two structurally different ends, and form the "core" of the actin cytoskeleton. An actin cytoskeleton is thus a three dimensional structure that results from the interaction between actin filaments and other molecules that associate with the actin filaments (e.g., cross-linking proteins such as filamin). The actin cytoskeleton mediates a variety of biological functions in all eukaryotic cells. In addition to providing a structural framework

around which cell shape and polarity are defined. its dynamic properties provide the driving force for cells to move and to divide.

According to the present invention, it has been discovered that agents which disrupt actin cytoskeletal organization control endothelial cell Nitric Oxide Synthase activity. In particular, agents that disrupt actin cytoskeletal organization upregulate endothelial cell Nitric Oxide Synthase activity.

According to the present invention, "agents that disrupt actin cytoskeletal organization" are compounds, natural or synthetic, that interfere with actin cytoskeletal organization. Typically such agents will interfere, for example, with stress fiber formation (contractile bundles of actin filaments and myosin), and/or focal contact (or *adhesion plaque*) assembly and upregulate endothelial cell Nitric Oxide Synthase activity. The effects of such agents in a cell or in a tissue on actin cytoskeletal organization can be measured according to any art recognized method. For example, a direct measure would be to perform phalloidin staining (Sigma) on intact cells. A person of ordinary skill in the art could then determine (and quantitate) the effects of the agents of the invention by examining, for example, the structure of the stained actin stress-fibers and comparing such structure with the one which is normal and characteristic of an untreated cell.

Agents that disrupt actin cytoskeletal organization can exert their effects at different levels and thus comprise different categories of agents useful for practicing the present invention. The different categories include agents from those that disrupt actin cytoskeletal organization at the nucleic acid level to agents that disrupt actin cytoskeletal organization at the protein level.

Agents that disrupt actin cytoskeletal organization at the nucleotide level include chemicals, antisense nucleic acids, antibodies, catalytic nucleic acids including ribozymes, and proteins which interfere with the expression of a gene that encodes a polypeptide which is a component of the actin cytoskeleton. Such exemplary polypeptides include but are not limited to actin, myosin, tropomyosin, troponin, titin, nebulin,  $\alpha$ -actinin, myomesin, C protein, filamin, talin, vinculin, capping protein, fibronectin receptor, ezrin, radixin, moiesin and the like.

Agents that disrupt actin cytoskeletal organization at the protein level include organic molecules that inhibit or alter the formation and organization of the actin cytoskeleton by interfering (e.g., via antibody binding, etc.) or altering (e.g., via post-translational modification) an individual component of the actin cytoskeleton. Specifically included are proteins, peptides and lipid derivatives. Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology.

-20-

Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W.R. (1986) The Experimental Foundations of Modern Immunology Wiley & Sons, Inc., New York; Roitt, I. (1991) Essential Immunology, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F(ab')<sub>2</sub> fragment, retains both of the antigen binding sites of an intact antibody. Similarly, an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment, retains one of the antigen binding sites of an intact antibody molecule. Proceeding further, Fab fragments consist of a covalently bound antibody light chain and a portion of the antibody heavy chain denoted Fd. The Fd fragments are the major determinant of antibody specificity (a single Fd fragment may be associated with up to ten different light chains without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the epitope of the antigen, and framework regions (FRs), which maintain the tertiary structure of the paratope (see, in general, Clark, 1986; Roitt, 1991). In both the heavy chain Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) separated respectively by three complementarity determining regions (CDR1 through CDR3). The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely responsible for antibody specificity.

It is now well-established in the art that the non-CDR regions of a mammalian antibody may be replaced with similar regions of conspecific or heterospecific antibodies while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently joined to human FR and/or Fc/pFc' regions to produce a functional antibody. Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of humanized murine RSV antibodies in which at least a portion of the murine FR regions have been replaced by FR regions of human origin. Such antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

Thus, as will be apparent to one of ordinary skill in the art, the present invention also

-21-

provides for F(ab')<sub>2</sub>, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric F(ab')<sub>2</sub> fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric Fab fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced by homologous human or non-human sequences. The present invention also includes so-called single chain antibodies.

10 In certain embodiments, agents that disrupt actin cytoskeletal organization include myosin light chain kinase (MLCK- Ser/Thr kinases) inhibitors, myosin light chain phosphatase (MLCP) stimulators, protein kinase N (PKN) inhibitors, phosphatidylinositol 4-phosphate 5-kinase (PIP5K) inhibitors, and cytochalasin D. Exemplary myosin light chain kinase inhibitors include BDM [2,3-butanedione 2-monoxime], ML-7 [1-(5-iodonaphthalene-  
15 1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride], ML-9 [1-(5-chloronaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride], wortmannin, H-7 [1-(5-isoquinoline  
sulphonyl)-2-methylpiperazine dihydro-chloride], Fasudil (HA1077) [Hexahydro-1-(5-isoquinolinesulphonyl)-1H-1,4- diazepine], W-7 [N-(6-Aminohexyl)-5-chloro-1-naphthalenesulfonamide] and A-3 [N-(6-Aminoethyl)-5-chloro-1-naphthalenesulfonamide].

20 In preferred embodiments, agents that disrupt actin cytoskeletal organization include BDM, ML-7, H-7 and cytochalasin D. Exemplary PKN inhibitors include "dominant negative" PKN peptides and purine analogues such as 6-thioguanine. Exemplary PIP5K inhibitors include "dominant negative" PIP5K peptides. Exemplary MLCP stimulators include nucleic acids that encode functional MLCP proteins and peptides (i.e., maintain the phosphatase activity of MLCP)  
25 and that are overexpressed (via an expression vector) in the cells of interest of a subject according to the invention.using genetic approaches well known in the art. Cytochalasin D is a preferred agent of the invention that belongs to the family of mold metabolites called cytochalasins. Cytochalasin D is thought to exert its function as an agent that disrupts actin cytoskeletal organization by affecting actin polymerization. Other members of the cytochalasin family share  
30 this property (e.g., Cytochalasin B), and are thus useful according to the invention.

Examples of agents that disrupt actin cytoskeletal organization also include "dominant negative" polypeptides of the polypeptide components of the actin cytoskeleton, some of which

-22-

are exemplified above. A dominant negative polypeptide is an inactive variant of a protein, which, by interacting with the cellular machinery, displaces an active protein from its interaction with the cellular machinery or competes with the active protein, thereby reducing the effect of the active protein. For example, a dominant negative receptor which binds a ligand but does not  
5 transmit a signal in response to binding of the ligand can reduce the biological effect of expression of the ligand. Likewise, a dominant negative catalytically-inactive kinase which interacts normally with target proteins but does not phosphorylate the target proteins can reduce phosphorylation of the target proteins in response to a cellular signal. Similarly, a dominant negative transcription factor which binds to a promoter site in the control region of a gene but  
10 does not increase gene transcription can reduce the effect of a normal transcription factor by occupying promoter binding sites without increasing transcription.

The end result of the application of or expression of a dominant negative polypeptide is a reduction in function of active proteins. One of ordinary skill in the art can assess the potential for a dominant negative variant of a protein, and using standard mutagenesis techniques to create  
15 one or more dominant negative variant polypeptides. For example, given the teachings contained herein and in the art, one of ordinary skill in the art can modify the sequence of a polypeptide (or the gene encoding a polypeptide) of an actin cytoskeletal component (as described earlier, e.g., actin, myosin, filamin, etc.) by site-specific mutagenesis, scanning mutagenesis, partial gene deletion or truncation, and the like. See, e.g., U.S. Patent No. 5,580,723 and Sambrook et al.,  
20 *Molecular Cloning: A Laboratory Manual*. Second Edition, Cold Spring Harbor Laboratory Press, 1989. The skilled artisan then can test the population of mutagenized polypeptides for diminution in a selected activity (e.g., impaired myosin light chain phosphorylation and upregulation of eNOS activity) and/or for retention of such an activity. Other similar methods for creating and testing dominant negative variants of a protein will be apparent to one of ordinary  
25 skill in the art.

Other examples of agents that disrupt actin cytoskeletal organization include polypeptides which bind to components of the actin cytoskeleton and to complexes of the components of the actin cytoskeleton and binding partners. The invention, therefore, embraces peptide binding agents which, for example, can be antibodies or fragments of antibodies having the ability to  
30 selectively bind to components of the actin cytoskeleton. Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology.

A *rho* GTPase is a small, membrane-bound, Ras-related GTP-binding protein that



-23-

functions by binding and hydrolyzing GTP. *Rho* GTPases function as molecular switches, cycling between an inactive GDP-bound conformation and an active GTP-bound conformation. According to the present invention, "*rho* GTPase function inhibitors" are compounds, natural or synthetic, that inhibit the normal function and localization of *rho* GTPases (i.e., impair GTP binding by *rho* GTPases) and upregulate endothelial cell Nitric Oxide Synthase activity. Such compounds can inhibit *rho* GTPase function at different levels and thus comprise different categories of agents useful for practicing the present invention. The different categories include agents from those that inhibit *rho* GTPases at the nucleic acid level to agents that inhibit *rho* GTPases at the protein level.

10 Agents that inhibit *rho* GTPases at the nucleotide level include chemicals, antisense nucleic acids, antibodies, catalytic nucleic acids including ribozymes, and proteins which repress expression of a *rho* GTPase gene locus.

Agents that inhibit *rho* GTPases at the protein level include organic molecules that alter the intrinsic GTPase activity of the *rho* GTP-binding protein, organic molecules that inhibit GDP/GTP exchange, and organic molecules that inhibit or alter post-translational modifications of *rho* GTPases. Specifically included are proteins, peptides and lipid derivatives.

Examples of agents that inhibit or reduce the intrinsic GTPase activity of a *rho* GTP-binding protein include cyclosporin, and "dominant negative" polypeptides of the *rho* GTPase. A dominant negative polypeptide is as described previously.

20 Dominant negative *rho* GTPase proteins include variants in which a portion of the GTP catalytic site has been mutated or deleted to reduce or eliminate GTP binding. Other examples include *rho* GTPase variants in which the conserved CAAX motif at their carboxy-terminus has been mutated or deleted to reduce or eliminate post-translational modification. (C, cysteine; A, aliphatic amino acid; X, any amino acid). One of ordinary skill in the art can readily prepare such modifications. Examples of dominant negative *rho* GTPase peptides are described in the Examples section and include N19RhoA and CAAXRhoA.

Other examples of agents that inhibit or reduce the intrinsic GTPase activity of a *rho* GTP-binding protein include polypeptides which bind to *rho* GTPase polypeptides and to complexes of *rho* GTPase polypeptides and binding partners. The invention, therefore, embraces peptide binding agents which, for example, can be antibodies or fragments of antibodies having the ability to selectively bind to *rho* GTPase polypeptides. Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology.

30

Examples of agents that inhibit the GDP/GTP exchange include proteins and peptides that inhibit GDP-dissociation such as Ly-GDI and RhoGDI-3. Preferably, using genetic approaches well known in the art, such proteins and peptides can be overexpressed (via an expression vector) in the cells of interest of a subject according to the invention.

5        Post-translational modifications of *rho* GTPases are important in that they are necessary for the proper attachment (and thus function) of the *rho* GTPases to the cell membrane. If *rho* GTPase polypeptides cannot be properly modified (or if they are overmodified), they accumulate in the cytosol and are rendered inactive. Examples of agents that inhibit post-translational modifications of *rho* GTPases include geranylgeranylation inhibitors and guanine nucleotide  
10    exchange inhibitors.

Geranylgeranylation inhibitors are compounds (natural or synthetic) that interfere with the geranylgeranylation of *rho* GTPases, and include proteins, peptides and lipid derivatives. Thus, geranylgeranylation inhibition of *rho* GTPases can occur either by preventing geranylgeranyl-pyrophosphate synthesis, or by inhibiting the enzyme geranylgeranyl transferase  
15    (GGT) which attaches geranylgeranyl-pyrophosphate to the CAAX motif of *rho* GTPases. Geranylgeranyl-pyrophosphate synthesis inhibition can be performed by preventing or inhibiting the formation of any of the intermediates in the geranylgeranyl-pyrophosphate synthesis pathway. Examples include mevalonate inhibitors, isopentenyl-pyrophosphate inhibitors, geranyl-pyrophosphate inhibitors, farnesyl-pyrophosphate inhibitors and geranylgeranyl-pyrophosphate  
20    inhibitors. Examples of such compounds include farnesyl-transferase inhibitors disclosed in U.S. patents 5,705,686 and 5,602,098, inhibitors of geranylgeranyl-transferase disclosed in U.S. patent 5,470,832, the disclosure of which is incorporated herein by reference, and a-hydroxyfarnesylphosphonic acid. Additional geranylgeranyl-transferase inhibitors include GGTI-298 (Finder, JD et al., *J Biol Chem*, 1997, 272:13484-13488).

25        Guanine nucleotide exchange inhibitors are agents that also post-translationally modify and inactivate *rho* GTPases. They include bacterial protein toxins that ADP-ribosylate or glucosylate *rho* GTPases, or compounds that inhibit *rho* GTPase-specific guanine nucleotide exchange factor (GEF). Preferred such agents according to the invention include *Clostridium botulinum* C3 transferase. The C3 transferase enzymatically catalyses the transfer of ADP from  
30    NADH to Asp-41 of *rho*, enhancing the conversion of active *rho* GTPase (GTP) to inactive *rho* GTPase (*rho* GDP) (See the Examples section also). The C3 transferase is administered in protein form, or more preferably, its cDNA is expressed using an expression vector in the cells

-25-

of interest of a subject according to the invention. *Rho* GTPase-specific guanine nucleotide exchange factor inhibitors include chemicals, antisense nucleic acids, antibodies, catalytic nucleic acids including ribozymes, proteins which repress expression of a *rho* GTPase-specific guanine nucleotide exchange factor gene locus, proteins, peptides (including dominant-negative peptides and antibodies), and the like.

According to the invention, agents that disrupt actin cytoskeletal organization are used excluding *rho* GTPase function inhibitors as agents useful in upregulating eNOS activity. The invention can involve use of a *rho* GTPase function inhibitor (including a HMG-CoA reductase inhibitor), however, only if used together with an agent that disrupts actin cytoskeletal organization other than a *rho* GTPase function inhibitor.

HMG-CoA reductase inhibitors inhibit post-translational modifications of *rho* GTPases by preventing mevalonate synthesis and consequently geranylgeranylpyrophosphate synthesis, an isoprenoid that is attached to the CAAX motif of *rho* GTPases. Examples of HMG-CoA reductase inhibitors include some which are commercially available, such as simvastatin (U.S. Patent No. 4, 444,784), lovastatin (U.S. Patent No. 4,231,938), pravastatin sodium (U.S. Patent No. 4,346,227), fluvastatin (U.S. Patent No. 4,739,073), atorvastatin (U.S. Patent No. 5,273,995), cerivastatin, and numerous others described in U.S. Patent No. 5,622,985, U.S. Patent No. 5,135,935, U.S. Patent No. 5,356,896, U.S. Patent No. 4,920,109, U.S. Patent No. 5,286,895, U.S. Patent No. 5,262,435, U.S. Patent No. 5,260,332, U.S. Patent No. 5,317,031, U.S. Patent No. 5,283,256, U.S. Patent No. 5,256,689, U.S. Patent No. 5,182,298, U.S. Patent No. 5,369,125, U.S. Patent No. 5,302,604, U.S. Patent No. 5,166,171, U.S. Patent No. 5,202,327, U.S. Patent No. 5,276,021, U.S. Patent No. 5,196,440, U.S. Patent No. 5,091,386, U.S. Patent No. 5,091,378, U.S. Patent No. 4,904,646, U.S. Patent No. 5,385,932, U.S. Patent No. 5,250,435, U.S. Patent No. 5,132,312, U.S. Patent No. 5,130,306, U.S. Patent No. 5,116,870, U.S. Patent No. 5,112,857, U.S. Patent No. 5,102,911, U.S. Patent No. 5,098,931, U.S. Patent No. 5,081,136, U.S. Patent No. 5,025,000, U.S. Patent No. 5,021,453, U.S. Patent No. 5,017,716, U.S. Patent No. 5,001,144, U.S. Patent No. 5,001,128, U.S. Patent No. 4,997,837, U.S. Patent No. 4,996,234, U.S. Patent No. 4,994,494, U.S. Patent No. 4,992,429, U.S. Patent No. 4,970,231, U.S. Patent No. 4,968,693, U.S. Patent No. 4,963,538, U.S. Patent No. 4,957,940, U.S. Patent No. 4,950,675, U.S. Patent No. 4,946,864, U.S. Patent No. 4,946,860, U.S. Patent No. 4,940,800, U.S. Patent No. 4,940,727, U.S. Patent No. 4,939,143, U.S. Patent No. 4,929,620, U.S. Patent No. 4,923,861, U.S. Patent No. 4,906,657, U.S. Patent No. 4,906,624 and U.S. Patent No. 4,897,402, the disclosures of which

-26-

patents are incorporated herein by reference.

Other *rho* GTPase function inhibitors not described in the above categories and useful according to the invention include agents that inhibit *rho* GTPase activation via a receptor-mediated signaling pathway. Such agents include protein kinase C inhibitors, Gq protein inhibitors (e.g., C-terminal antibodies, dominant-negative Gq mutants, etc.), tyrosine kinase inhibitors (e.g., genistein, etc.), tyrosine phosphatase stimulators, *rho* GTPase-activating protein stimulators, inhibitors of integrins and adhesion molecules, adapter protein (Shc and Sos) inhibitors, inhibitors of growth factor receptors (bradykinin receptor, EGF receptor, FGF receptor, etc.), and inhibitors of proteins containing Pleckstrin homology domains which bind G-protein- $\beta\gamma$ , but only where administration of these agents is effective to increase ecNOS activity.

According to another aspect of the invention, a method of screening for identifying an agent that disrupts actin cytoskeletal organization for treatment of subjects who would benefit from increased endothelial cell Nitric Oxide Synthase activity in a tissue, is provided. The method involves identifying an agent that disrupts actin cytoskeletal organization suspected of increasing endothelial cell Nitric Oxide Synthase activity, and determining whether or not the agent that disrupts actin cytoskeletal organization produces an increase in endothelial cell Nitric Oxide Synthase activity *in vivo* or *in vitro*. Agents that disrupt actin cytoskeletal organization according to this invention can be identified by confirming that the inhibitor produces increased ecNOS activity in a model system compared to a control, using any of the model systems described herein, and also inhibits at least one other actin cytoskeletal disruptive function as determined in any of the model systems described herein and/or other model systems known in the art.

The invention also involves the co-administration of agents that are not agents that disrupt actin cytoskeletal organization but that can act cooperatively, additively or synergistically with such agents that disrupt actin cytoskeletal organization to increase ecNOS activity. Thus, ecNOS substrates which are converted by ecNOS to nitric oxide and cofactors enhancing such conversion, can be co-administered with the agents that disrupt actin cytoskeletal organization according to the invention. Such ecNOS substrates (e.g. L-arginine) and cofactors (e.g., NADPH, tetrahydrobiopterin, etc.) may be natural or synthetic.

The invention also involves the co-administration of agents that are not agents that disrupt actin cytoskeletal organization but that can act cooperatively, additively or synergistically with such agents that disrupt actin cytoskeletal organization to increase ecNOS activity. Thus, ecNOS substrates which are converted by ecNOS to nitric oxide can be co-administered with the agents

-27-

that disrupt actin cytoskeletal organization according to the invention. Such ecNOS substrates may be natural or synthetic, although the preferred substrate is L-arginine.

Likewise, there are other agents besides agents that disrupt actin cytoskeletal organization, that are not substrates of ecNOS, and that can increase ecNOS activity. Agents belonging to these categories are not therefore agents that disrupt actin cytoskeletal organization as described herein, and can be used in co-administrations with agents that disrupt actin cytoskeletal organization in cocktails. Examples of categories of such agents are estrogens and ACE inhibitors. Estrogens are a well defined category of molecules known by those of ordinary skill in the art, and will not be elaborated upon further herein. All share a high degree of structural similarity. ACE inhibitors also have been well characterized, although they do not always share structural homology.

Angiotensin converting enzyme, or ACE, is an enzyme which catalyzes the conversion of angiotensin I to angiotensin II. ACE inhibitors include amino acids and derivatives thereof, peptides, including di and tri peptides and antibodies to ACE which intervene in the renin-angiotensin system by inhibiting the activity of ACE thereby reducing or eliminating the formation of pressor substance angiotensin II. ACE inhibitors have been used medically to treat hypertension, congestive heart failure, myocardial infarction and renal disease. Classes of compounds known to be useful as ACE inhibitors include acylmercapto and mercaptoalkanoyl prolines such as captopril (US Patent Number 4,105,776) and zofenopril (US Patent Number 4,316,906), carboxyalkyl dipeptides such as enalapril (US Patent Number 4,374,829), lisinopril (US Patent Number 4,374,829), quinapril (US Patent Number 4,344,949), ramipril (US Patent Number 4,587,258), and perindopril (US Patent Number 4,508,729), carboxyalkyl dipeptide mimics such as cilazapril (US Patent Number 4,512,924) and benazapril (US Patent Number 4,410,520), phosphinylalkanoyl prolines such as fosinopril (US Patent Number 4,337,201) andtrandolopril.

This invention also contemplates co-administration of agents that increase the production of NO by ecNOS without affecting ecNOS expression, as do ACE inhibitors or administration of ecNOS substrate and/or ecNOS cofactors. Estrogens upregulate Nitric Oxide Synthase expression whereas ACE inhibitors do not affect expression, but instead influence the efficiency of the action of Nitric Oxide Synthase on L-arginine. Thus, activity can be increased in a variety of ways. In general, activity is increased by the reductase inhibitors of the invention by increasing the amount of the active enzyme present in a cell versus the amount present in a cell

absent treatment with the reductase inhibitors according to the invention.

The invention also involves the co-administration of "at least one different agent that disrupts actin cytoskeletal organization" (second agent that disrupts actin cytoskeletal organization) that can act cooperatively, additively or synergistically with a first agent that  
5 disrupts actin cytoskeletal organization of the invention to increase ecNOS activity. Thus, "at least one different agent that disrupts actin cytoskeletal organization" is meant to include one or more agent(s) that disrupts actin cytoskeletal organization that is (are) different to the first agent that disrupts actin cytoskeletal organization of the invention and can include a HMG-CoA reductase inhibitor and/or a *rho* GTPase function inhibitor. In one embodiment, when the agent  
10 that disrupts actin cytoskeletal organization according to the invention is co-administered in combination with "at least one different agent that disrupts actin cytoskeletal organization" and the "at least one different agent that disrupts actin cytoskeletal organization" is a HMG-CoA reductase inhibitor, the subject is nonhypercholesterolemic.

The agents that disrupt actin cytoskeletal organization are administered in effective  
15 amounts. In general, an effective amount is any amount that can cause an increase in Nitric Oxide Synthase activity in a desired cell or tissue, and preferably in an amount sufficient to cause a favorable phenotypic change in a condition such as a lessening, alleviation or elimination of a symptom or of a condition.

In general, an effective amount is that amount of a pharmaceutical preparation that alone,  
20 or together with further doses or co-administration of other agents, produces the desired response. This may involve only slowing the progression of the disease temporarily, although more preferably, it involves halting the progression of the disease permanently or delaying the onset of or preventing the disease or condition from occurring. This can be monitored by routine methods. Generally, doses of active compounds would be from about 0.01 mg/kg per day to 1000  
25 mg/kg per day. It is expected that doses ranging from 50-500 mg/kg will be suitable, preferably orally and in one or several administrations per day.

Such amounts will depend, of course, on the particular condition being treated, the severity of the condition, the individual patient parameters including age, physical condition, size and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific  
30 route of administration and like factors within the knowledge and expertise of the health practitioner. Lower doses will result from certain forms of administration, such as intravenous administration. In the event that a response in a subject is insufficient at the initial doses applied,

-29-

higher doses (or effectively higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits. Multiple doses per day are contemplated to achieve appropriate systemic levels of compounds. It is preferred generally that a maximum dose be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however, that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reasons.

The agents that disrupt actin cytoskeletal organization useful according to the invention may be combined, optionally, with a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid or liquid fillers, diluents or encapsulating substances which are suitable for administration into a human. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being co-mingled with the molecules of the present invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficacy.

The pharmaceutical compositions may contain suitable buffering agents, including: acetic acid in a salt; citric acid in a salt; boric acid in a salt; and phosphoric acid in a salt.

The pharmaceutical compositions also may contain, optionally, suitable preservatives, such as: benzalkonium chloride; chlorobutanol; parabens and thimerosal.

A variety of administration routes are available. The particular mode selected will depend, of course, upon the particular drug selected, the severity of the condition being treated and the dosage required for therapeutic efficacy. The methods of the invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of the active compounds without causing clinically unacceptable adverse effects. Such modes of administration include oral, rectal, topical, nasal, interdermal, or parenteral routes. The term "parenteral" includes subcutaneous, intravenous, intramuscular, or infusion. Intravenous or intramuscular routes are not particularly suitable for long-term therapy and prophylaxis.

The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well-known in the art of pharmacy. All methods include the step of bringing the active agent into association with a carrier which constitutes one or more

-30-

accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product.

Compositions suitable for oral administration may be presented as discrete units, such as capsules, tablets, lozenges, each containing a predetermined amount of the active compound. Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir or an emulsion.

Compositions suitable for parenteral administration conveniently comprise a sterile aqueous preparation of reductase inhibitors, which is preferably isotonic with the blood of the recipient. This aqueous preparation may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation also may be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid may be used in the preparation of injectables. Carrier formulation suitable for oral, subcutaneous, intravenous, intramuscular, etc. administrations can be found in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, PA.

Other delivery systems can include time-release, delayed release or sustained release delivery systems. Such systems can avoid repeated administrations of the active compound, increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer base systems such as poly(lactide-glycolide), copolyoxalates, polycaprolactones, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and polyanhydrides. Microcapsules of the foregoing polymers containing drugs are described in, for example, U.S. Patent 5,075,109. Delivery systems also include non-polymer systems that are: lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-di-and tri-glycerides; hydrogel release systems; sylastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which the active compound is contained



-31-

in a form within a matrix such as those described in U.S. Patent Nos. 4,452,775, 4,675,189, and 5,736,152, and (b) diffusional systems in which an active component permeates at a controlled rate from a polymer such as described in U.S. Patent Nos. 3,854,480, 5,133,974 and 5,407,686. In addition, pump-based hardware delivery systems can be used, some of which are adapted for  
5 implantation.

Use of a long-term sustained release implant may be desirable. Long-term release, as used herein, means that the implant is constructed and arranged to deliver therapeutic levels of the active ingredient for at least 30 days, and preferably 60 days. Long-term sustained release implants are well-known to those of ordinary skill in the art and include some of the release  
10 systems described above.

According to another aspect of the invention, a method for increasing blood flow in a tissue of a subject is provided. The method involves administering to a subject in need of such treatment a first agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the tissue of the subject, provided that  
15 the first agent is not a *rho* GTPase function inhibitor, H7, or fasudil. Fasudil (a substituted isoquinolinesulfonyl compound- also known as HA1077), described in U.S. Patent No. 4,678,783 as a compound with vasodilating properties, was not known to act as an agent that disrupts actin cytoskeletal organization resulting in increased ecNOS expression prior to the present invention.

In important embodiments a second agent is co-administered to a subject with a condition  
20 treatable by the second agent in an amount effective to treat the condition, whereby the delivery of the second agent to a tissue of the subject is enhanced as a result of the increased blood flow from administering the first agent of the invention (an agent that disrupts actin cytoskeletal organization). The "second agent" may be any pharmacological compound or diagnostic agent, as desired. Preferred second agents are agents having a site of action in the brain. Such agents  
25 include analeptic, analgetic, anesthetic, adrenergic agent, anti-adrenergic agent, amino acids, antagonists, antidote, anti-anxiety agent, anticholinergic, anticolvunsant, antidepressant, anti-emetic, anti-epileptic, antihypertensive, antifibrinolytic, antihyperlipidemia, antimigraine, antinauseant, antineoplastic (brain cancer), antiobsessional agent, antiparkinsonian, antipsychotic, appetite suppressant, blood glucose regulator, cognition adjuvant, cognition enhancer,  
30 dopaminergic agent, emetic, free oxygen radical scavenger, glucocorticoid, hypocholesterolemic, hyperlipidemic, histamine H2 receptor antagonists, immunosuppressant, inhibitor, memory adjuvant, mental performance enhancer, mood regulator, mydriatic,

-32-

neuromuscular blocking agent, neuroprotective, NMDA antagonist, post-stroke and post-head trauma treatment, psychotropic, sedative, sedative-hypnotic, serotonin inhibitor, tranquilizer, and treatment of cerebral ischemia, calcium channel blockers, free radical scavengers - antioxidants, GABA agonists, glutamate antagonists, AMPA antagonists, kainate antagonists, competitive and non-competitive NMDA antagonists, growth factors, opioid antagonists, phosphatidylcholine precursors, serotonin agonists, sodium- and calcium-channel blockers, and potassium channel openers.

In addition to the foregoing brain-specific categories of agents, examples of categories of other pharmaceutical agents that can be used as second agents include: adrenergic agent; adrenocortical steroid; adrenocortical suppressant; alcohol deterrent; aldosterone antagonist; amino acid; ammonia detoxicant; anabolic; analeptic; analgesic; androgen; anesthesia, adjunct to; anesthetic; anorectic; antagonist; anterior pituitary suppressant; anthelmintic; anti-acne agent; anti-adrenergic; anti-allergic; anti-amebic; anti-androgen; anti-anemic; anti-anginal; anti-anxiety; anti-arthritic; anti-asthmatic; anti-atherosclerotic; antibacterial; anticholelithic; anticholelithogenic; anticholinergic; anticoagulant; anticoccidal; anticonvulsant; antidepressant; antidiabetic; antidiarrheal; antidiuretic; antidote; anti-emetic; anti-epileptic; anti-estrogen; antifibrinolytic; antifungal; antiglaucoma agent; antihemophilic; antihemorrhagic; antihistamine; antihyperlipidemia; antihyperlipoproteinemic; antihypertensive; anti-infective; anti-infective, topical; anti-inflammatory; antikeratinizing agent; antimalarial; antimicrobial; antimigraine; antimitotic; antimycotic; antinauseant; antineoplastic; antineutropenic; antiobessional agent; antiparasitic; antiparkinsonian; antiperistaltic; antipneumocystic; antiproliferative; antiprostatic hypertrophy; antiprotozoal; antipruritic; antipsychotic; antirheumatic; antischistosomal; antiseborrheic; antisecretory; antispasmodic; antithrombotic; antitussive; anti-ulcerative; anti-urolithic; antiviral; appetite suppressant; benign prostatic hyperplasia therapy agent; blood glucose regulator; bone resorption inhibitor; bronchodilator; carbonic anhydrase inhibitor; cardiac depressant; cardioprotectant; cardiotonic; cardiovascular agent; choleric; cholinergic; cholinergic agonist; cholinesterase deactivator; coccidiostat; cognition adjuvant; cognition enhancer; depressant; diagnostic aid; diuretic; dopaminergic agent; ectoparasiticide; emetic; enzyme inhibitor; estrogen; fibrinolytic; fluorescent agent; free oxygen radical scavenger; gastrointestinal motility effector; glucocorticoid; gonad-stimulating principle; hair growth stimulant; hemostatic; histamine H2 receptor antagonists; hormone; hypocholesterolemic; hypoglycemic; hypolipidemic; hypotensive; imaging agent;

-33-

immunizing agent; immunomodulator; immunoregulator; immunostimulant; immunosuppressant; impotence therapy adjunct; inhibitor; keratolytic; LNRH agonist; liver disorder treatment; luteolysin; memory adjuvant; mental performance enhancer; mood regulator; mucolytic; mucosal protective agent; mydriatic; nasal decongestant; neuromuscular blocking agent; neuroprotective; NMDA antagonist; non-hormonal sterol derivative; oxytocic; plasminogen activator; platelet activating factor antagonist; platelet aggregation inhibitor; post-stroke and post-head trauma treatment; potentiator; progestin; prostaglandin; prostate growth inhibitor; prothyrotropin; psychotropic; pulmonary surface; radioactive agent; regulator; relaxant; repartitioning agent; scabicide; sclerosing agent; sedative; sedative-hypnotic; selective adenosine A1 antagonist; serotonin antagonist; serotonin inhibitor; serotonin receptor antagonist; steroid; stimulant; suppressant; symptomatic multiple sclerosis; synergist; thyroid hormone; thyroid inhibitor; thyromimetic; tranquilizer; treatment of amyotrophic lateral sclerosis; treatment of cerebral ischemia; treatment of Paget's disease; treatment of unstable angina; uricosuric; vasoconstrictor; vasodilator; vulnerary; wound healing agent; xanthine oxidase inhibitor.

In another aspect of the invention, the agent that disrupts actin cytoskeletal organization is "co-administered," which means administered substantially simultaneously with another agent. By substantially simultaneously, it is meant that the agent that disrupts actin cytoskeletal organization is administered to the subject close enough in time with the administration of the other agent, whereby the two compounds may exert an additive or even synergistic effect, i.e. on increasing eNOS activity or on delivering a second agent to a tissue via increased blood flow.

### Examples

*"Upregulation of endothelial cell Nitric Oxide Synthase by HMG CoA Reductase Inhibitors"*

#### Experimental Procedures

All standard culture reagents were obtained from JRH Bioscience (Lenexa, KS). Unless indicated otherwise, all reagents were purchased from Sigma Chemical Co. (St. Louis, MO). [ $\alpha$ - $^{32}$ P]CTP (3000 Ci/mmol) was supplied by New England Nuclear. Purified human LDL was obtained from Calbiochem (San Diego, CA; lot#730793) and Biomedical Technologies Inc. (Stoughton, MA; lot#9030197). The level of endotoxin was determined by the chromogenic Limulus amebocyte assay (BioWhittaker Inc., Walkersville, MD). The antibody detection kit (Enhanced Chemiluminescence) and the nylon nucleic acid (Hybond) and protein (PVDF)

transfer membranes were purchased from Amersham Corp. (Arlington Heights, IL). Simvastatin and lovastatin were obtained from Merck, Sharp, and Dohme, Inc. (West Point, PA). Since endothelial cells lack lactonases to process simvastatin and lovastatin to their active forms, these HMG-CoA reductase inhibitors were chemically activated prior to their use with methods well known in the art and as previously described (Laufs, U et al., *J Biol Chem*, 1997, 272:31725-31729).

#### Cell Culture:

Human endothelial cells were harvested from saphenous veins and cultured as described (15). For transfection studies, bovine aortic endothelial cells of less than 3 passages were cultured in a growth medium containing DMEM (Dulbecco's Modified Eagle's Medium), 5 mmol/L L-glutamine (Gibco), and 10% fetal calf serum (Hyclone Lot#1114577). For all experiments, the endothelial cells were placed in 10% lipoprotein-deficient serum (Sigma, Lot#26H94031) for 48 h prior to treatment conditions. In the indicated experiments, endothelial cells were pretreated with actinomycin D (5 mg/ml) for 1 h prior to treatment with ox-LDL and/or simvastatin. Cellular viability as determined by cell count, morphology, and Trypan blue exclusion was maintained for all treatment conditions.

#### Preparation of LDL:

The LDL was prepared by discontinuous ultracentrifugation according to the method of Chung et al. with some modification (*Methods Enzymol*, 1984, 128:181-209). Fresh plasma from a single donor was anticoagulated with heparin and filtered through a Sephadex G-25 column equilibrated with PBS. The density was adjusted to 1.21 g/ml by addition of KBr (0.3265 g/ml plasma). A discontinuous NaCl/KBr gradient was established in Beckman Quick-Seal centrifuge tubes (5.0 ml capacity) by layering 1.5 ml of density-adjusted plasma under 3.5 ml of 0.154 M NaCl in Chelex-100-treated water (BioRad, Hercules, CA). After ultracentrifugation at 443,000 x g and 7°C for 45 min in a Beckman Near Vertical Tube 90 rotor (Beckman L8-80M ultracentrifuge), the yellow band in the upper middle of the tube corresponding to LDL was removed by puncturing with a needle and withdrawing into a syringe. The KBr was removed from the LDL by dialyzing with three changes of sterile PBS, pH 7.4, containing 100 µg/ml polymyxin B.

The purity of the LDL samples was confirmed by SDS/polyacrylamide and cellulose

-35-

acetate gel electrophoresis. Cholesterol and triglyceride content were determined as previously described (Liao, JK et al., *J Biol Chem*, 1995, 270:319-324.). The LDL protein concentration was determined by the method of Lowry et al., (*J Biol Chem*, 1951, 193:265-275.). For comparison, commercially-available LDL (Biomedical Technologies Inc., Stoughton, MA; Calbiochem, San Diego, CA) were characterized and used in selected experiments.

#### Oxidation of LDL:

Oxidized LDL was prepared by exposing freshly-isolated LDL to  $\text{CuSO}_4$  (5-10 mM) at 37°C for various duration (6-24 h). The reaction was stopped by dialyzing with three changes of sterile buffer (150  $\mu\text{mol/L}$  NaCl, 0.01% EDTA and 100  $\mu\text{g/ml}$  polymyxin B, pH 7.4) at 4°C. The degree of LDL oxidation was estimated by measuring the amounts of thiobarbituric acid reactive substances (TBARS) produced using a fluorescent assay for malondialdehyde as previously described (Yagi, KA, *Biochem Med*, 1976, 15:212-21.). The extent of LDL modification was expressed as nanomoles of malondialdehyde per mg of LDL protein. Only mild to moderate oxidized LDL with TBARS values between 12 and 16 nmol/mg LDL protein (i.e. 3 to 4 nmol/mg LDL cholesterol) were used in this study. All oxidatively-modified LDL samples were used within 24 h of preparation.

#### Northern Blotting:

Equal amounts of total RNA (10-20 mg) were separated by 1.2% formaldehyde-agarose gel electrophoresis and transferred overnight onto Hybond nylon membranes. Radiolabeling of human full-length ecNOS cDNA (Verbeuren, TJ et al., *Circ Res*, 1986, 58:552-564, Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666) was performed using random hexamer priming, [ $^{32}\text{P}$ ]CTP, and Klenow (Pharmacia). The membranes were hybridized with the probes overnight at 45°C in a solution containing 50% formamide, 5 X SSC, 2.5 X Denhardt's Solution, 25 mM sodium phosphate buffer (pH 6.5), 0.1% SDS, and 250 mg/ml salmon sperm DNA. All Northern blots were subjected to stringent washing conditions (0.2 X SSC/0.1% SDS at 65°C) prior to autoradiography. RNA loading was determined by rehybridization with human GAPDH probe.

#### Western Blotting:

Cellular proteins were prepared and separated on SDS/PAGE as described (Liao, JK et al., *J Biol Chem*, 1995, 270:319-324). Immunoblotting was performed using a murine monoclonal

-36-

antibody to human ecNOS (1:400 dilution, Transduction Laboratories, Lexington, KY). Immunodetection was accomplished using a sheep anti-mouse secondary antibody (1:4000 dilution) and the enhanced chemiluminescence (ECL) kit (Amersham Corp., Arlington Heights, IL). Autoradiography was performed at 23°C and the appropriate exposures were quantitated by densitometry.

#### Assay for ecNOS Activity:

The ecNOS activity was determined by a modified nitrite assay as previously described (Misko, TP et al., *Analytical Biochemistry*, 1993, 214:11-16, Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666). Briefly, endothelial cells were treated for 24 h with ox-LDL in the presence and absence of simvastatin (0.1 to 1 mM). After treatment, the medium was removed, and the cells were washed and incubated for 24 h in phenol red-free medium. After 24 h, 300 µl of conditioned medium was mixed with 30 µl of freshly prepared 2,3-diaminonaphthalene (1.5 mmol/L DAN in 1 mol/L HCl). The mixture was protected from light and incubated at 20°C for 10 min. The reaction was terminated with 15 µl of 2.8 mol/L NaOH. Fluorescence of 1-(H)-naphthotriazole was measured with excitation and emission wavelengths of 365 and 450 nm, respectively. Standard curves were constructed with known amounts of sodium nitrite. Nonspecific fluorescence was determined in the presence of LNMA (5 mmol/L).

#### Nuclear Run-on Assay:

Confluent endothelial cells (~5 x 10<sup>7</sup> cells) grown in LPDS were treated with simvastatin (1 mM) or 95%O<sub>2</sub> for 24 h. Nuclei were isolated and in vitro transcription was performed as previously described (Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666). Equal amounts (1 mg) of purified, denatured full-length human ecNOS, human b-tubulin (ATCC #37855), and linearized pGEM-3z cDNA were vacuum-transferred onto nitrocellulose membranes using a slot blot apparatus (Schleicher & Schuell). Hybridization of radiolabeled mRNA transcripts to the nitrocellulose membranes was carried out at 45°C for 48 h in a buffer containing 50% formamide, 5 X SSC, 2.5 X Denhardt's solution, 25 mM sodium phosphate buffer (pH 6.5), 0.1% SDS, and 250 mg/ml salmon sperm DNA. The membranes were then washed with 1 x SSC/0.1% SDS for 1 h at 65°C prior to autoradiography for 72 h at -80°C.

#### Transfection Assays:

-37-

For transient transfections, bovine rather than human endothelial cells were used because of their higher transfection efficiency by the calcium-phosphate precipitation method (12% vs <4%) (Graham, FL and Van der Erb, AJ. *Virology*, 1973, 52:456-457). We used the human ecNOS promoter construct, F1.LUC, which contains a -1.6 kb 5'-upstream sequence linked to the luciferase reporter gene as described by Zhang et al. (*J Biol Chem*, 1995, 270:15320-15326). Bovine endothelial cells (60%-70% confluent) were transfected with 30 mg of the indicated constructs: p.LUC (no promoter), pSV2.LUC (SV40 early promoter), or F1.LUC. As an internal control for transfection efficiency, pCMV.bGal plasmid (10 mg) was co-transfected in all experiments. Preliminary results using b-galactosidase staining indicate that cellular transfection efficiency was approximately 10% to 14%.

Endothelial cells were placed in lipoprotein-deficient serum for 48 h after transfection and treated with ox-LDL (50 mg/ml, TBARS 12.4 nmol/mg) in the presence and absence of simvastatin (1 mM) for an additional 24 h. The luciferase and b-galactosidase activities were determined by a chemiluminescence assay (Dual-Light, Tropix, Bedford, MA) using a Berthold L9501 luminometer. The relative promoter activity was calculated as the ratio of luciferase to b-galactosidase activity. Each experiment was performed three times in triplicate.

#### Data Analysis:

Band intensities were analyzed densitometrically by the National Institutes of Health Image program (Rasband, W, NIH Image program, v 1.49, National Institutes of Health, Bethesda, 1993). All values are expressed as mean  $\pm$  SEM compared to controls and among separate experiments. Paired and unpaired Student's t tests were employed to determine any significant changes in densitometric values, nitrite production, and promoter activities. A significant difference was taken for P values less than 0.05.

#### Example 1: Cell Culture

Relatively pure (>95%) human endothelial cell cultures were confirmed by their morphological features (i.e. cuboidal, cobble-stone, contact inhibited) using phase-contrast microscopy and by immunofluorescent staining with anti-Factor VIII antibodies (Gerson, RJ et al., *Am J Med*, 1989, 87:28-38). For all experimental conditions, there were no observable adverse effects of ox-LDL or HMG-CoA reductase inhibitors on cellular morphology, cell number, immunofluorescent staining, and Trypan blue exclusion (>95%). Higher concentrations of ox-

-38-

LDL (>100 mg/ml) with greater oxidative modification (i.e. TBARS values of >30 nmol/mg) caused vacuolization and some detachment of endothelial cells after 24 h. Neither simvastatin (0.01 to 0.1 mmol/L) nor lovastatin (10 mmol/L) produced any noticeable adverse effects on human endothelial cell for up to 96 h. However, higher concentrations of simvastatin (>15 mmol/L) or lovastatin (>50 mmol/L) caused cytotoxicity after 36 h, and therefore, were not used.

#### Example 2: Characterization of LDL

SDS/polyacrylamide gel electrophoresis of native or unmodified LDL revealed a single band (~510 kD) corresponding to ApoB-100 (data not shown). Similarly, cellulose acetate electrophoresis revealed only one band corresponding to the presence of a single class of low-density lipids (density of 1.02 to 1.06 g/ml). The LDL had a protein, cholesterol, and triglyceride concentration of  $6.3 \pm 0.2$ ,  $2.5 \pm 0.1$ , and  $0.5 \pm 0.1$  mg/ml, respectively. In contrast, lipoprotein-deficient serum was devoid of both apoB-100 protein and low-density lipid bands, and had non-detectable levels of cholesterol. There was no detectable level of endotoxin (<0.10 EU/ml) in the lipoprotein-deficient serum or ox-LDL samples by the chromogenic Limulus amebocyte assay.

In addition, there was no apparent difference between our own preparation and commercially-obtained LDL samples in terms of electrophoretic mobility. Native LDL had a TBARS value of  $0.3 \pm 0.2$  nmol/mg, but after exposure to human saphenous vein endothelial cells in lipoprotein-deficient media for 72 h, this value increased to  $3.1 \pm 0.4$  nmol/mg. Copper-oxidized LDL had TBARS values ranging from  $4.6 \pm 0.5$  to  $33.1 \pm 5.2$  nmol/mg. The degree of ox-LDL used in this study was mild to moderate with TBARS value ranging from 12 to 16 nmol/mg LDL protein (i.e. 3 to 4 nmol/mg LDL cholesterol).

#### Example 3: Effect of ox-LDL and HMG-CoA Reductase Inhibitors on ecNOS Protein

We have previously shown that ox-LDL (50 mg/ml) downregulates ecNOS expression (Liao, JK et al., *J Biol Chem*, 1995, 270:319-324). Compared to untreated cells, treatment with ox-LDL (50 mg/ml, TBARS 12.2 nmol/mg) caused a  $54\% \pm 6\%$  decrease in ecNOS protein after 48 h ( $p < 0.01$ ,  $n=4$ ). There was no difference between our preparation of ox-LDL and commercially-available ox-LDL with similar TBARS values in terms of the degree of ecNOS downregulation. Addition of simvastatin (0.01 mmol/L) did not significantly affect the downregulation of ecNOS protein by ox-LDL ( $57\% \pm 8\%$  decrease,  $p > 0.05$ ,  $n=4$ ). However, in the presence of 0.1 mmol/L of simvastatin, ox-LDL no longer produce any significant decrease



in ecNOS protein levels ( $4\% \pm 7\%$  decrease,  $p < 0.01$ ,  $n=4$ ). Higher concentrations of simvastatin (1 and 10 mmol/L) resulted in not only a reversal of ecNOS downregulation by ox-LDL, but also significant increases in ecNOS protein levels above baseline ( $146\% \pm 9\%$  and  $210\% \pm 12\%$ , respectively,  $p < 0.05$ ,  $n=4$ ). Simvastatin or lovastatin (10 mmol/L) which were not chemically-activated had no effect on ecNOS expression.

In a time-dependent manner, treatment with ox-LDL (50 mg/ml, TBARS 12.2 nmol/mg) decreased ecNOS protein expression by 34% (5%, 67% (8% and 86 (5% after 24 h, 72 h, and 96 h, respectively ( $p < 0.05$  for all values,  $n=4$ )). Compared to ox-LDL alone, co-treatment with simvastatin (0.1 mmol/L) attenuated the decrease in ecNOS protein level after 24 h (15% (2% vs 34% (5%,  $p < 0.05$ ,  $n=4$ ). Longer incubation with simvastatin (0.1 mmol/L) for 72 h and 96 h not only reversed ox-LDL's inhibitory effects on ecNOS expression, but also increased ecNOS protein levels by 110% (6% and 124% (6% above basal expression ( $p < 0.05$ ,  $n=4$ ). Thus, compared to ox-LDL alone, co-treatment with simvastatin produced a 1.3-, 3.3- and 8.9-fold increase ecNOS protein levels after 24 h, 72 h, and 96 h, respectively.

#### Example 4: Effect of ox-LDL and HMG-CoA Reductase Inhibitors on ecNOS mRNA

The effect of simvastatin on ecNOS mRNA levels occurred in a time-dependent manner and correlated with its effect on ecNOS protein levels. Northern analyses showed that ox-LDL (50 mg/ml, TBARS 15.1 nmol/mg) produced a time-dependent  $65 \pm 5\%$  and  $91 \pm 4\%$  decrease in ecNOS mRNA levels after 48 h and 72 h, respectively ( $p < 0.01$ ,  $n=3$ ). Compared to ox-LDL at the indicated time points, co-treatment with simvastatin 0.1 mmol/L increased ecNOS mRNA levels by 6.3-fold after 48 h and 14.5-fold after 72 h ( $p < 0.01$  for all values,  $n=3$ ).

To determine whether treatment with another HMG-CoA reductase inhibitor have similar effect as simvastatin, we treated endothelial cells with lovastatin. Again, ox-LDL decreased steady-state ecNOS mRNA by  $52 \pm 5\%$  after 24 h ( $p < 0.01$ ,  $n=3$ ). Treatment with lovastatin (10 mmol/L) not only reversed the inhibitory effects of ox-LDL on ecNOS mRNA, but also caused a  $40 \pm 9\%$  increase in ecNOS mRNA level compared to that of untreated cells. Compared to ox-LDL alone, co-treatment with lovastatin caused a 3.6-fold increase in ecNOS mRNA levels after 24 h. Treatment with lovastatin alone, however, produced 36% increase in ecNOS mRNA levels compared to untreated cells ( $p < 0.05$ ,  $n=3$ ).

#### Example 5: Effect of ox-LDL and Simvastatin on ecNOS Activity

-40-

The activity of ecNOS was assessed by measuring the LNMA-inhibitable nitrite production from human endothelial cells (Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666). Basal ecNOS activity was  $8.8 \pm 1.4$  nmol/500,000 cells/24 h. Treatment with ox-LDL (50 mg/ml, TBARS 16 nmol/mg) for 48 h decreased ecNOS-dependent nitrite production by  $94 \pm 3\%$  ( $0.6 \pm 0.5$  nmol/500,000 cells/24 h,  $p < 0.001$ ). Co-treatment with simvastatin (0.1 mmol/L) significantly attenuated this downregulation resulting in a  $28 \pm 3\%$  decrease in ecNOS activity compared to untreated cells ( $6.4 \pm 0.3$  nmol/500,000 cells/24 h,  $p < 0.05$ ). Co-treatment with a higher concentration of simvastatin (1 mmol/L) not only completely reversed the downregulation of ecNOS by ox-LDL, but also, resulted in a  $45 \pm 6\%$  increase in ecNOS activity compared to baseline ( $12.8 \pm 2.7$  nmol/500,000 cells/24 h,  $p < 0.05$ ).

#### Example 6: Effect of Simvastatin on ecNOS mRNA Stability

The post-transcriptional regulation of ecNOS mRNA was determined in the presence of the transcriptional inhibitor, actinomycin D (5 mg/ml). Oxidized LDL (50 mg/ml, TBARS 13.1 nmol/mg) shortened the half-life of ecNOS mRNA ( $t_{1/2}$   $35 \pm 3$  h to  $14 \pm 2$  h,  $p < 0.05$ ,  $n=3$ ). Co-treatment with simvastatin (0.1 mmol/L) prolonged the half-life of ecNOS mRNA by 1.6-fold ( $t_{1/2}$   $22 \pm 3$  h,  $p < 0.05$ ,  $n=3$ ). Treatment with simvastatin alone prolonged ecNOS mRNA half-life by 1.3-fold over baseline ( $t_{1/2}$   $43 \pm 4$  h,  $p < 0.05$ ,  $n=3$ ).

#### Example 7: Effect of Simvastatin on ecNOS Gene Transcription

To determine whether the effects of simvastatin on ecNOS expression occurs at the level of ecNOS gene transcription, we performed nuclear run-on assays using endothelial cells treated with simvastatin (1 mmol/L) for 24 h. Preliminary studies using different amounts of radiolabelled RNA transcripts demonstrate that under our experimental conditions, hybridization was linear and nonsaturable. The density of each ecNOS band was standardized to the density of its corresponding b-tubulin. The specificity of each band was determined by the lack of hybridization to the nonspecific pGEM cDNA vector. In untreated endothelial cells (control), there was constitutive ecNOS transcriptional activity (relative index of 1.0). Treatment with simvastatin (1 mmol/L) did not significantly affect ecNOS gene transcription compared to that of untreated cells (relative index of  $1.2 \pm 0.3$ ,  $p > 0.05$ ,  $n=4$ ). However, treatment of endothelial cells with hyperoxia (95%  $O_2$ ) significantly increased ecNOS gene expression (relative index of 2.5,  $p < 0.05$ ,  $n=4$ ).

-41-

To further confirm the effects of simvastatin on ecNOS gene transcription by a different method, we transfected bovine aortic endothelial cells using a -1600 to +22 nucleotide ecNOS 5'-promoter construct linked to a luciferase reporter gene (F1.LUC) (Zhang, R et al., *J Biol Chem*, 1995, 270:15320-15326). This promoter construct contains putative cis-acting elements for activator protein (AP)-1 and -2, sterol regulatory element-1, retinoblastoma control element, shear stress response element (SSRE), nuclear factor-1 (NF-1), and cAMP response element (CRE). Treatment with ox-LDL (50 mg/ml, TBARS 14.5 nmol/mg), simvastatin (1  $\mu$ mol/L), alone or in combination, did not significantly affect basal F1 promoter activity. However, laminar fluid shear-stress (12 dynes/cm<sup>2</sup> for 24 h) was able to induce F1 promoter activity by 16-fold after 24 h (data not shown) indicating that the F1 promoter construct is functionally-responsive if presented with the appropriate stimulus.

#### Example 8: Effect of Simvastatin and Lovastatin on ecNOS Expression

To further characterize the effects of HMG-CoA reductase inhibitors on the upregulation ecNOS expression, we treated endothelial cells with simvastatin (0.1 mmol/L) for various durations (0-84 h). Treatment with simvastatin (0.1 mmol/L) increased ecNOS protein levels by 4 ( 6%, 21 ( 9%, 80 ( 8%, 90 ( 12%, and 95 ( 16% after 12 h, 24 h, 48 h, 72 h, and 84 h, respectively ( $p < 0.05$  for all time points after 12 h,  $n=4$ ). Higher concentrations of simvastatin similarly increased ecNOS protein levels, but in significantly less time compared to lower concentrations of simvastatin (data not shown).

In a concentration-dependent manner, treatment with simvastatin (0.01 to 10 mmol/L, 48 h) increased ecNOS expression by 1 ( 6%, 80 ( 8%, 190 ( 10% and 310 ( 20%, respectively ( $p < 0.05$  for concentrations ( 0.1 mmol/L,  $n=4$ ). The upregulation of ecNOS expression by simvastatin, therefore, is dependent upon both the concentration and duration of simvastatin treatment. For comparison, treatment with lovastatin (0.1 to 10 mmol/L, 48 h) also increased ecNOS expression in a concentration-dependant manner (10 ( 6%, 105 ( 8% and 180 ( 11%, respectively,  $p < 0.05$  for concentrations  $> 0.1$  mmol/L,  $n=3$ ) but significantly less effectively than simvastatin at comparable concentrations. Therefore, at the same concentration, simvastatin had greater effects on ecNOS expression compared to lovastatin. These results are consistent with reported IC<sub>50</sub> values for simvastatin and lovastatin (4 nmol/L and 19 nmol/L, respectively) (Van Vliet, AK et al., *Biochem Pharmacol*, 1996, 52:1387-1392).

Example 9: Effect of L-Mevalonate on ecNOS Expression

To confirm that the effects of simvastatin on ecNOS expression were due to the inhibition of endothelial HMG CoA reductase, endothelial cells were treated with ox-LDL (50 mg/ml, TBARS 15.1 nmol/mg), simvastatin (1 mmol/L), alone or in combination, in the presence of L-mevalonate (100 mmol/L). Treatment with ox-LDL decreased ecNOS expression by 55% (6% after 48 h which was completely reversed and slightly upregulated in the presence of simvastatin (1 mmol/L) (150% (8% above basal expression) ( $p < 0.05$  for both,  $n=3$ ).

Compared to endothelial cells treated with ox-LDL and simvastatin, addition of L-mevalonate reduced ecNOS protein by  $50\% \pm 5\%$  ( $p < 0.05$ ,  $n=3$ ). Furthermore, the upregulation of ecNOS expression by simvastatin alone (2.9-fold increase,  $p < 0.05$ ,  $n=3$ ) was completely reversed by co-treatment with L-mevalonate. Treatment with L-mevalonate alone did not have any appreciable effects on basal ecNOS expression ( $p > 0.05$ ,  $n=3$ ). Similar findings were also observed with L-mevalonate and lovastatin.

*"HMG-CoA Reductase Inhibitors Reduce Cerebral Infarct Size by Upregulating endothelial cell Nitric Oxide Synthase"*

Experimental ProceduresCell Culture:

Human endothelial cells were harvested from saphenous veins using Type II collagenase (Worthington Biochemical Corp., Freehold, NJ) as previously described. Cells of less than three passages were grown to confluence in a culture medium containing Medium 199, 20 mM HEPES, 50 mg/ml ECGS (Collaborative Research Inc., Bedford, MA), 100 mg/ml heparin sulfate, 5 mM L-glutamine (Gibco), 5% fetal calf serum (Hyclone, Logan, UT), and antibiotic mixture of penicillin (100 U/ml)/ streptomycin (100 mg/ml)/Fungizone (1.25 mg/ml). For all experiments, the endothelial cells were grown to confluence before any treatment conditions. In some experiments, cells were pretreated with actinomycin D (5 mg/ml) for 1 h prior to treatment with HMG-CoA reductase inhibitors.

Exposure of Endothelial Cells to Hypoxia:

Confluent endothelial cells grown in 100 mm culture dishes were treated with HMG-CoA reductase inhibitors and then placed without culture dish covers in humidified airtight incubation

chambers (Billups-Rothenberg, Del Mar, CA). The chambers were gassed with 20% or 3% O<sub>2</sub>, 5% CO<sub>2</sub>, and balanced nitrogen for 10 min prior to sealing the chambers. The chambers were maintained in a 37°C incubator for various durations (0-48 h) and found to have less than 2% variation in O<sub>2</sub> concentration as previously described (Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666). Cellular confluence and viability were determined by cell count, morphology, and trypan blue exclusion.

#### In vitro Transcription Assay:

Confluent endothelial cells (5 x 10<sup>7</sup> cells) were treated with simvastatin (1 mM) in the presence of 20% or 3% O<sub>2</sub> for 24 h. Nuclei were isolated and in vitro transcription was performed as previously described (Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666). Equal amounts (1 mg) of purified, denatured full-length human ecNOS, human  $\beta$ -tubulin (ATCC #37855), and linearized pGEM-3z cDNA were vacuum-transferred onto nitrocellulose membranes using a slot blot apparatus (Schleicher & Schuell). Hybridization of radiolabeled mRNA transcripts to the nitrocellulose membranes was carried out at 45°C for 48 h in a buffer containing 50% formamide, 5 X SSC, 2.5 X Denhardt's solution, 25 mM sodium phosphate buffer (pH 6.5), 0.1% SDS, and 250 mg/ml salmon sperm DNA. The membranes were then washed with 1 x SSC/0.1% SDS for 1 h at 65°C prior to autoradiography for 72 h at -80°C. Band intensities were subjected to analyses by laser densitometry.

#### Assay for Nitrite Accumulation:

The amount of NO produced by ecNOS was determined by nitrite accumulation in the conditioned medium. Nitrite accumulation was determined by measuring the conversion of 2,3-diaminonaphthalene (1.5 mM of DAN in 1 M of HCl) and nitrite to 1-(H)-naphthotriazole as previously described (13,24). Nonspecific fluorescence was determined in the presence of LNMA (5 mM). Previous studies with nitrate reductase indicate that the nitrite to nitrate concentration in the medium was approximately 5:1 and that this ratio did not vary with exposure to 20% or 3% O<sub>2</sub> concentration.

#### Murine Model of Cerebral Vascular Ischemia:

Adult male (18-20 g) wildtype SV-129 mice (Taconic farm, Germantown, NY) and ecNOS mutant mice (Huang, PL et al., *Nature*, 1995, 377:239-242.) were subcutaneously-injected

with 0.2, 2, or 20 mg of activated simvastatin per kg body weight or saline (control) once daily for 14 days. Ischemia was produced by occluding the left middle cerebral artery (MCA) with a coated 8.0 nylon monofilament under anesthesia as described (Huang, Z et al., *J Cereb Blood Flow Metab*, 1996, 16:981-987, Huang, Z et al., *Science*, 1994, 265:1883-1885, Hara, H et al., *J Cereb Blood Flow Metab*, 1997, 1:515-526). Arterial blood pressure, heart rate, arterial oxygen pressure, and partial pressure of carbon dioxide were monitored as described (Huang, Z et al., *J Cereb Blood Flow Metab*, 1996, 16:981-987, Huang, Z et al., *Science*, 1994, 265:1883-1885, Hara, H et al., *J Cereb Blood Flow Metab*, 1997, 1:515-526). The filaments were withdrawn after 2 hours and after 24 h, mice were either sacrificed or tested for neurological deficits using a well-established, standardized, observer-blinded protocol as described (Huang, Z et al., *J Cereb Blood Flow Metab*, 1996, 16:981-987, Huang, Z et al., *Science*, 1994, 265:1883-1885, Hara, H et al., *J Cereb Blood Flow Metab*, 1997, 1:515-526). The motor deficit score range from 0 (no deficit) to 2 (complete deficit).

Brains were divided into five coronal 2-mm sections using a mouse brain matrix (RBM-200C, Activated Systems, Ann Arbor, MI, USA). Infarction volume was quantitated with an image analysis system (M4, St. Catharines, Ontario, Canada) on 2% 2,3,5-triphenyltetrazolium chloride stained 2-mm slices. The levels of serum cholesterol, creatinine and transaminases were determined by the Tufts University Veterinary Diagnostic Laboratory (Grafton, MA).

#### Assay for ecNOS Activity from Tissues:

The ecNOS activities in mice aortae and brains were measured by the conversion of [<sup>3</sup>H]arginine to [<sup>3</sup>H]citrulline in the presence and absence of LNMA (5 mM) as described earlier.

#### Quantitative Reverse Transcription-Polymerase Chain Reaction:

Total RNA from mouse aortae and brains was isolated by the guanidinium isothiocyanate method and reverse transcribed using oligo-dT (mRNA Preamplification reagents; Gibco BRL) and Taq polymerase (Perkin-Elmer). One tenth of the sDNA was used as template for the PCR reaction. Approximately 0.2 nmol of the following primers amplifying a 254-bp fragment of murine ecNOS cDNA were used: 5'Primer: 5'-GGGCTCCCTCCTTCCGGCTGCCACC-3' (SEQ ID NO. 1) and 3'Primer: 5'-GGATCCCTGGAAAAGGCGGTGAGG-3' (SEQ ID NO. 2) (Hara, H et al., *J Cereb Blood Flow Metab*, 1997, 1:515-526). For amplification of glyceraldehyde 3-phosphate dehydrogenase (GAPDH), 0.1 nmol of the following primers amplifying a 452-bp

-45-

fragment were used: 5'Primer: 5'-ACCACAGTCCATGCCATCAC-3' (SEQ ID NO. 3) and 3' Primer: 5'-TCCACCACCCTGTTGCTGTA-3'(SEQ ID NO. 4). Denaturing was performed at 94°C for 30 s, annealing at 60°C for 30 s, and elongation at 72°C for 60 s. Preliminary results indicated that the linear exponential phase for ecNOS and GAPDH polymerization was 30-35 cycles and 20-25 cycles, respectively.

#### Example 10: Cell Culture

Relatively pure (>98%) human saphenous vein endothelial cell cultures were confirmed by their morphological features (ie. cuboidal, cobble-stone, contact inhibited) using phase-contrast microscopy and immunofluorescent-staining with antibodies to Factor VIII. There were no observable adverse effects of HMG-CoA reductase inhibitors, L-mevalonic acid, or hypoxia on cellular morphology. However, higher concentrations of simvastatin (>15 mmol/L) or lovastatin (>50 mmol/L) caused cytotoxicity after 36 h, and therefore, were not used. Otherwise, cellular confluency and viability as determined by trypan blue exclusion were maintained for all treatment conditions described.

#### Example 11: Effects of HMG-CoA Reductase Inhibitors on ecNOS Activity

The activity of ecNOS was assessed by measuring the LNMA-inhibitable nitrite accumulation from human endothelial cells (Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666). The ratio of nitrite to nitrate production under our culture condition was approximately 5:1 and was similar for hypoxia and normoxia (data not shown). Basal ecNOS activity at 20% O<sub>2</sub> was  $6.0 \pm 3.3$  nmol/500,000 cells/24 h. Exposure of endothelial cells to 3% O<sub>2</sub> for 24 h decreased nitrite production by  $75 \pm 14\%$  ( $1.5 \pm 0.9$  nmol/500,000 cells/24 h,  $p < 0.01$ ). Treatment with simvastatin (1 mM) not only completely reversed the downregulation of ecNOS by hypoxia, but resulted in a 3-fold increase in ecNOS activity over basal activity ( $18 \pm 5.0$  nmol/500,000 cells/24 h,  $p < 0.05$ ). This upregulation of ecNOS activity was attenuated by the addition of L-mevalonate (400 mM) ( $9.6 \pm 1.3$  nmol/500,000 cells/24 h,  $p < 0.05$ ). Interestingly, simvastatin (1 mM) alone upregulated nitrite production 5-fold ( $30 \pm 6.5$  nmol/500,000 cells/24 h,  $p < 0.01$ ), which was completely blocked by L-mevalonate (400 mM) ( $8.6 \pm 2.9$  nmol/500,000 cells/24 h,  $p < 0.05$ ). Similar findings were observed with lovastatin, but at 10-fold higher concentration compared to that of simvastatin.

Example 12: Effects of HMG-CoA Reductase Inhibitors on ecNOS Protein and mRNA Levels

In a concentration-dependent manner, treatment with simvastatin (0.01 to 10 mM, 48 h) increased ecNOS expression by 1 (6%, 80 (8%, 190 (10% and 310 (20%, respectively ( $p < 0.05$  for concentrations (0.1 mM,  $n=4$ ). Treatment with simvastatin (0.1 mM) increased ecNOS protein levels in a time-dependent manner by 4 (6%, 21 (9%, 80 (8%, 90 (12%, and 95 (16% after 12 h, 24 h, 48 h, 72 h, and 84 h, respectively ( $p < 0.05$  for all time points after 12 h,  $n=4$ ) (data not shown). Another HMG-CoA reductase inhibitor, lovastatin, also increased ecNOS protein levels in a time-, and concentration-dependent manner (data not shown). Because lovastatin has a higher IC<sub>50</sub> value for HMG-CoA reductase compared to that of simvastatin, it was 10-fold less potent in upregulating ecNOS protein levels than simvastatin at equimolar concentrations.

We have previously shown that hypoxia downregulates ecNOS protein expression (Liao, JK et al., *J Clin Invest*, 1995, 96:2661-2666). Compared to normoxia (20% O<sub>2</sub>), exposure to hypoxia (3% O<sub>2</sub>) resulted in a  $46 \pm 4$  % and  $75 \pm 3$  % reduction in ecNOS protein levels after 24 h and 48 h, respectively ( $p < 0.01$ ,  $n=3$ ). In a concentration-dependent manner, treatment with simvastatin produced a progressive reversal of hypoxia-mediated downregulation of ecNOS protein levels after 48 h. At higher concentrations of simvastatin (1 and 10 mM), ecNOS protein levels were upregulated to  $159 \pm 13$  % and  $223 \pm 21$  % of basal levels ( $p < 0.05$ ,  $n=3$ ). Co-treatment with L-mevalonic acid (400 mM) completely blocked simvastatin-induced increase in ecNOS protein levels after 48 h ( $35 \pm 2.4$  %). Treatment with L-mevalonic acid alone, however, did not produce any significant effects on basal ecNOS protein levels in untreated cells exposed to hypoxia ( $25 \pm 3.9$  %,  $p > 0.05$ ,  $n=3$ ). In addition, simvastatin which was not chemically-activated had no effect on ecNOS expression. These results indicate that simvastatin- and lovastatin-mediated increases in ecNOS protein expression are mediated by inhibition of endothelial HMG-CoA reductase.

To determine whether changes in ecNOS protein levels are due to changes in ecNOS steady-state mRNA levels, we performed Northern blotting on endothelial cells exposed to normoxia and hypoxia in the presence or absence of simvastatin (1 mM) and lovastatin (10  $\mu$ M). Simvastatin alone increased ecNOS mRNA levels to  $340 \pm 24$  % ( $p < 0.01$ ,  $n=3$ ). Exposure of endothelial cells to hypoxia reduced ecNOS mRNA levels by  $70\% \pm 2$  % and  $88 \pm 4$  % after 24 h and 48 h with respect to GAPDH mRNA levels, respectively. Co-treatment with simvastatin not only completely reversed hypoxia-mediated decrease in ecNOS mRNA levels, but increased



-47-

ecNOS mRNA levels to  $195 \pm 12\%$  and  $530 \pm 30\%$  of basal levels after 24 h and 48 h, respectively ( $p < 0.01$ ,  $n=3$ ). Similarly, lovastatin ( $10 \mu\text{M}$ ) alone increased ecNOS message to  $350 \pm 27\%$  under hypoxia and  $410 \pm 21\%$  alone ( $p < 0.01$ ,  $n=3$ ). Neither simvastatin nor lovastatin caused any significant change in G-protein  $\alpha$  and  $\beta$ -actin mRNA levels under normoxic or hypoxic conditions. These results indicate that the effects of HMG-CoA reductase inhibitors are relatively selective in terms of their effects on ecNOS mRNA expression.

#### Example 13: Effects of HMG-CoA Reductase Inhibitors on ecNOS mRNA Half-life

The half-life of ecNOS mRNA was determined in the presence of actinomycin D ( $5 \text{ mg/ml}$ ). Hypoxia shortened the half-life of ecNOS mRNA from  $28 \pm 4 \text{ h}$  to  $13 \pm 3 \text{ h}$ . Treatment with simvastatin ( $1 \text{ mM}$ ) increased ecNOS half-life to  $46 \pm 4 \text{ h}$  and  $38 \pm 4 \text{ h}$  under normoxic and hypoxic conditions, respectively ( $p < 0.05$  for both,  $n=3$ ). These results suggest that HMG-CoA reductase inhibitors prevent hypoxia-mediated decrease in ecNOS expression by stabilizing ecNOS mRNA.

#### Example 14: Effects of HMG-CoA Reductase Inhibitors on ecNOS Gene Transcription

Nuclear run-on assays showed that hypoxia caused a  $85 \pm 8\%$  decrease in ecNOS gene transcription ( $p < 0.01$ ,  $n=3$ ). Treatment with simvastatin ( $1 \text{ mM}$ ) did not produce any significant affect on hypoxia-mediated decrease in ecNOS gene transcription ( $83 \pm 6\%$  decrease in ecNOS gene transcription,  $p > 0.05$  compared to hypoxia alone). Furthermore, simvastatin alone produced minimal increase in ecNOS gene transcription under normoxic condition ( $20 \pm 5\%$  increase in ecNOS gene transcription,  $p < 0.05$  compared to normoxia control).

Preliminary studies using different amounts of radiolabeled RNA transcripts demonstrate that under our experimental conditions, hybridization was linear and nonsaturable. The density of each ecNOS band was standardized to the density of its corresponding ( $\beta$ -tubulin band, relative intensity). To exclude the possibility that changes in ( $\beta$ -tubulin gene transcription are caused by hypoxia or simvastatin, another gene, GAPDH, was included on each of the nuclear run-on blots. Similar relative indices were obtained when ecNOS gene transcription was standardized to GAPDH gene transcription. The specificity of each band was determined by the lack of hybridization to the nonspecific pGEM cDNA vector.

#### Example 15: Effect of HMG-CoA Reductase Inhibitors on Mouse Physiology

-48-

To determine whether the upregulation of ecNOS by HMG-CoA reductase inhibitors occurs in vivo, SV-129 wild-type and ecNOS knockout mice were treated with 2 mg/kg simvastatin or saline subcutaneously for 14 days (n=8). The mean arterial blood pressures of wild-type and ecNOS mutant mice were as reported previously (Huang, PL et al., *Nature*, 1995, 377:239-242). The ecNOS mutants were relatively hypertensive. There was no significant change in mean arterial blood pressures of wild-type mice after 14 days of simvastatin treatment ( $81 \pm 7$  mmHg vs.  $93 \pm 10$  mmHg,  $p > 0.05$ ). There was also no significant group difference in heart rate, arterial blood gases and temporalis muscle temperature before ischemia or after reperfusion. Furthermore, there was no significant difference in the levels of serum cholesterol (control:  $147 \pm 10$  vs. simvastatin  $161 \pm 5.2$  mg/dl), creatinine and transaminases after treatment with simvastatin compared to control values.

Example 16: Effect of HMG-CoA Reductase Inhibitors on ecNOS Expression and Function in Mouse Aorta

The activity of ecNOS in the aortae of simvastatin-treated (2 mg/kg) and saline-injected mice was determined by measuring the LNMA-inhibitable conversion of arginine to citrulline (Figure 1 A). The ecNOS activity in aortae from simvastatin-treated mice was significantly higher than in the control group ( $0.39 \pm 0.09$  vs.  $0.18 \pm 0.04$  U/mg protein, n=8,  $p < 0.05$ ).

The ecNOS mRNA expression in the aortae of simvastatin-treated and -untreated mice was examined by quantitative RT-PCR (Figure 1 B). There was a significantly dose-dependent 3-fold increase of ecNOS message compared to that of GAPDH in simvastatin-treated mice (n=3). These findings indicate that simvastatin upregulates ecNOS expression and activity in vivo.

Example 17: Effect of HMG-CoA Reductase Inhibitors on Cerebral Ischemia in Mice

Endothelium-derived NO protects against ischemic cerebral injury (Huang, Z et al., *J Cereb Blood Flow Metab*, 1996, 16:981-987). Therefore we examined, whether the observed upregulation of ecNOS by simvastatin in vivo has beneficial effects on cerebral infarct size. Following treatment for 14 days with 2 mg/kg of simvastatin, cerebral ischemia was produced by occluding the left middle cerebral artery for 2 hours. After 22 hours of reperfusion, mice were tested for neurological deficits using a well-established, standardized, observer-blinded protocol. The neurological motor deficit score improved in simvastatin-treated mice (n=18) by almost 2-

fold compared to that of controls (n=12) ( $0.8 \pm 0.2$  vs.  $1.7 \pm 0.2$ ,  $p < 0.01$ ).

Simvastatin-treated wild-type mice (n=18) had 25% smaller cerebral infarct sizes compared to untreated animals ( $73.8 \pm 8.5$  mm<sup>3</sup> vs.  $100.7 \pm 7.3$  mm<sup>3</sup>, n=12,  $p < 0.05$ ). This effect was concentration-dependent (0.2, 2, 20 mg/kg simvastatin), persisted for up to 3 days, and also occurred with lovastatin treatment, albeit at higher relative concentrations (data not shown). Furthermore, simvastatin increase cerebral blood flow by 23% and 35% over basal values at concentrations of 2 mg/kg and 20 mg/kg, respectively (n=8,  $p < 0.05$  for both). These findings suggest, that simvastatin decreases cerebral infarct size and neurological deficits.

Finally, to demonstrate that the reduction of cerebral infarct sizes by simvastatin is due to the upregulation of eNOS, cerebral ischemia was applied to eNOS mutant mice lacking eNOS gene in the presence and absence of simvastatin (2 mg/kg, 14 days). There was no significant difference between the cerebral infarct sizes of simvastatin-treated and -untreated eNOS mutant mice (n=6,  $p < 0.05$ ). These findings indicate that the upregulation of eNOS mediates the beneficial effects of HMG-CoA reductase inhibitors on cerebral infarct size.

#### Example 18: Effect of HMG-CoA Reductase Inhibitors on eNOS Expression in Mouse Brain

The eNOS mRNA expression in the ischemic and contralateral (non-ischemic) hemispheres of mouse brain was examined by quantitative RT-PCR (Figure 2) with respect to GAPDH mRNA levels. Simvastatin-treated mice (n=3) (2 mg/kg, 14 days) showed a 1.5- to 2-fold increase in eNOS expression in the infarcted, ipsilateral hemisphere compared to the contralateral, non-infarcted side. In contrast, there was no difference in eNOS expression in untreated mice between their infarcted and non-infarcted hemispheres. These findings suggest that simvastatin may reduced cerebral infarct size by selectively increasing eNOS expression in the ischemic and hypoxic infarct zone.

#### *"Regulation of Endothelial Nitric Oxide Synthase Expression by Rho GTPases"*

#### Experimental Procedures

##### Materials:

Mevastatin, farnesylpyrophosphate, geranylgeranylpyrophosphate, and L-mevalonate were purchased from Sigma Chemical Corp. (St. Louis, MO). Mevastatin (compactin- a HMG-CoA reductase inhibitor) was chemically activated by alkaline hydrolysis prior to use as previously

-50-

described (Laufs, U et al., *J Biol Chem*, 1997, 272:31725-31729). FPT inhibitor I and -  
hydroxyfarnesylphosphonic acid were purchased from Calbiochem Corp. (La Jolla, CA). [ $\alpha$ -  
 $^{32}$ P]CTP (3000 Ci/mmol) and [ $^{35}$ S]GTP S (1250 Ci/mmol) were supplied by New England  
Nuclear. The antibody detection kit (Enhanced Chemiluminescence) and the nylon nucleic acid  
5 (Hybond) and protein (PVDF) transfer membranes were purchased from Amersham Corp.  
(Arlington Heights, IL). The *Clostridium botulinum* C3 transferase was purchased from List  
Biological Laboratories, Inc. (Campbell, CA). Recombinant *Escherichia coli* cytotoxic  
necrotizing factor (CNF)-1 and RhoA mutants were kindly provided by K. Aktories (University  
of Freiberg, Germany) and W. Moolenaar (Netherlands Cancer Institute, Netherlands),  
10 respectively.

#### Cell Culture

Human endothelial cells were harvested using Type II collagenase (Worthington  
Biochemical Corp., Freehold, NJ) as previously described (Laufs, U et al., *J Biol Chem*, 1997,  
15 272:31725-31729; Liao, JK et al., *J Biol Chem*, 1995, 270:319-324). Cells of less than three  
passages were grown in a culture medium containing Medium 199, 20 mM HEPES, 50 mg/ml  
ECGS (Collaborative Research Inc., Bedford, MA), 100 mg/ml heparin sulfate, 5 mM L-  
glutamine (Gibco), 5% fetal calf serum (Hyclone, Logan, UT), and antibiotic mixture of penicillin  
(100 U/ml)/ streptomycin (100 mg/ml)/Fungizone (1.25 mg/ml). Confluent endothelial cells were  
20 used for all treatment conditions. For transfection studies, bovine aortic endothelial cells of less  
than 3 passages were cultured in a growth medium containing DMEM (Dulbecco's Modified  
Eagle's Medium), 5 mM of L-glutamine (Gibco), and 10% fetal calf serum. Cellular viability was  
determined by cell count, morphology, and trypan blue exclusion.

#### Preparation of LDL

The LDL was prepared as described earlier. The extent of LDL oxidation was estimated  
by assaying for thiobarbituric acid reactive substances (TBARS) and expressed as nanomoles of  
malondialdehyde per mg of LDL protein, as described earlier. Only freshly-isolated LDL with  
TBARS values of less than 0.5 nmol/mg was used in this study.

#### Western Blotting

Proteins were prepared and separated on SDS/PAGE as described earlier. Immuno-

-51-

blotting was performed using monoclonal antibodies to ecNOS (1:400 dilution, Transduction Laboratories, Lexington, KY), to RhoA and RhoB (1:250 dilution, Santa Cruz Biotechnology Inc., Santa Cruz, CA), and to c-myc-tag (9E10, 1:200 dilution, Santa Cruz Biotechnology Inc.). Immunodetection was accomplished using a sheep anti-mouse secondary antibody (1:4000 dilution) or donkey anti-rabbit secondary antibody (1:4000 dilution) and the enhanced chemiluminescence (ECL) kit (Amersham Corp., Arlington Heights, IL). Autoradiography was performed as described earlier.

#### Assay for Rho GTP-binding Activity

The Rho GTP-binding activity was determined by immunoprecipitation of [<sup>35</sup>S]GTP S-labeled Rho. Briefly, membrane and cytosolic proteins were isolated as previously described (Liao, JK and Homcy, CJ, *J Clin Invest*, 1993, 92:2168-2172). Proteins (20 mg) from control and treated endothelial cells were incubated for 30 min at 37°C in a buffer containing [<sup>35</sup>S]GTP S (20 nM), GTP (2 mM), MgCl<sub>2</sub> (5 mM), EGTA (0.1 mM), NaCl (50 mM), creatinine phosphate (4 mM), phosphocreatinine kinase (5 units), ATP (0.1 mM), dithiothreitol (1 mM), leupeptin (100 mg/ml), aprotinin (50 mg/ml), and phenylmethanesulfonyl fluoride (PMSF, 2 mM). The assay was terminated with excess unlabeled GTP S (100 mM).

Samples were then resuspended in 100 ml of immunoprecipitation buffer containing Triton-X (1%), SDS (0.1%), NaCl (150 mM), EDTA (5 mM), Tris-HCl (25 mM, pH 7.4), leupeptin (10 mg/ml), aprotinin (10 mg/ml), and PMSF (2 mM). The RhoA or RhoB antisera were added to the mixture at a final dilution of 1:75. The samples were allowed to incubate for 16 h at 4°C with gentle mixing. The antibody-G-protein complexes were then incubated with 50 ml of protein A-Sepharose (1 mg/ml, Pharmacia Biotech Inc.) for 2 h at 4°C, and the immunoprecipitate was collected by centrifugation at 12,000 x g for 10 min.

Preliminary studies using Western analysis of the supernatant indicated that both RhoA and RhoB were completely immunoprecipitated under these conditions. The pellets were washed four times in a buffer containing HEPES (50 mM, pH 7.4), NaF (100 mM), sodium phosphate (50 mM), NaCl (100 mM), Triton X-100 (1%), and SDS (0.1%). The final pellet containing the immunoprecipitated [<sup>35</sup>S]GTP S-labeled Rho proteins was counted in a liquid scintillation counter (LS 1800, Beckman Instruments, Inc. Fullerton, CA). Nonspecific activity was determined in the presence of excess unlabeled GTP S (100 mM).

### Overexpression of Rho Mutants

For transfection studies, bovine rather than human endothelial cells were used because of their higher transfection efficiency by the calcium-phosphate precipitation method (12% vs <4%) (15). Bovine endothelial cells (60-70% confluent) were transfected with 15 mg of the indicated cDNAs: the insertless vector (pcDNA3), pcDNA3-c-myc-wtRhoA (wildtype RhoA), and pcDNA3-c-myc-N19RhoA (dominant-negative RhoA mutant) (Gebbink, M et al., *J Cell Biol*, 1997, 137:1603-1613). As an internal control for transfection efficiency, pCMV. b-Gal plasmid (5 mg) was co-transfected. Preliminary results using b-galactosidase staining indicate that cellular transfection efficiency was approximately 10% to 14%. The b-galactosidase activity was determined by a chemiluminescence assay (Dual-Light, Tropix, Bedford, MA) using a Berthold L9501 luminometer. Approximately 24 h after transfection, cells were harvested for immunoblot analyses of ecNOS expression. The ecNOS protein levels were then standardized to the corresponding levels of transfected RhoA expression as determined by antisera to the corresponding c-myc tag.

### Assay for ecNOS Activity

The ecNOS activity was determined by a modified nitrite assay as previously described. Briefly, endothelial cells grown in phenol-free medium were exposed to C3 transferase (50 mg/ml), FPP (10 mM), GGPP (5 mM), CNF-1 (200 ng/ml), or mevastatin (10 mM). After 24 h, conditioned medium (300 ml) was mixed with 30 ml of freshly-prepared 2,3-diaminonaphthalene (1.5 mM of DAN in 1 M of HCl). The mixture was protected from light and incubated at 20°C for 10 min. The reaction was terminated with 15 ml of 2.8 M of NaOH. Fluorescence of 1-(H)-naphthotriazole was measured with excitation and emission wavelengths of 365 and 450 nm, respectively. Standard curves were constructed with known amounts of sodium nitrite. Nonspecific fluorescence was determined in the presence of LNMA (3 mM). Previous studies with nitrate reductase indicate that the nitrite to nitrate concentration in the medium was approximately 5:1 and that this ratio did not vary under the described treatment conditions (Laufs, U et al., *J Biol Chem*, 1997, 272:31725-31729).

### Data Analysis

Band intensities from Northern and Western blots were analyzed as described earlier. Paired and unpaired Student's *t*-tests were employed to determine the significance of changes in

densitometric measurements, GTP-binding activities, and nitrite levels. A significant difference was taken for  $p < 0.05$ .

#### Example 19: Cell Culture

5           Relatively pure (>98%) human saphenous vein endothelial cell cultures were confirmed by their morphological features (i.e. cuboidal, cobble-stone, contact inhibited) using phase-contrast microscopy and immunofluorescent-staining with antibodies to Factor VIII (data not shown). There were no observable adverse effects of mevastatin, FPP, GGPP, C3 transferase, and CNF-1 on cellular viability. However, higher concentrations of mevastatin (>50 mM) or  
10       CNF-1 (>5 mg/ml) did produce cytotoxicity and therefore were not used. Cellular confluency and viability as determined by light microscopy and trypan blue exclusion were maintained for all treatment conditions described.

#### Example 20: Effects of isoprenoid intermediates on ecNOS mRNA expression

15           We have shown that inhibition of endothelial HMG-CoA reductase by lovastatin or simvastatin upregulates ecNOS expression and activity via increases in ecNOS mRNA stability (Examples 3-8 and Laufs, U et al., *J Biol Chem*, 1997, 272:31725-31729). Similarly, treatment of endothelial cells with mevastatin (10 mM) increased ecNOS steady-state mRNA levels by  $405 \pm 15\%$  after 24 h (Figure 3A). On a molar basis, we find that mevastatin is equally potent  
20       compared with lovastatin but approximately ten times less potent compared to simvastatin. This is consistent with their relative  $IC_{50}$  values for HMG-CoA reductase inhibition (Blum, CB. *Am. J. Cardiol*, 1994, 73:3D-11D).

          To determine which downstream isoprenoid intermediate in the cholesterol biosynthetic pathway regulates ecNOS expression, endothelial cells were treated with mevastatin (10 mM) in  
25       the presence or absence of isoprenoid intermediates, geranylgeranylpyrophosphate (GGPP) or farnesylpyrophosphate (FPP). Co-treatment with FPP (10 mM) mildly reduced ecNOS mRNA levels compared to mevastatin alone. However, co-treatment with GGPP (10 mM) completely reversed the upregulation of ecNOS mRNA levels by mevastatin. In a concentration-dependent manner, GGPP reversed the effects of mevastatin (10 mM) with complete reversal occurring at  
30       a GGPP concentration of 5 mM (Figure 3B). Interestingly, treatment with GGPP (10 mM) alone did not significantly affect basal ecNOS mRNA levels.

          Similarly, treatment with mevastatin (10 mM) increased ecNOS protein levels by  $180 \pm$

-54-

11% after 24 h ( $p < .05$ ,  $n = 4$ ) (Figure 4). Co-treatment with FPP (10 mM) or LDL (1 mg/ml) did not significantly reverse the effects of mevastatin on ecNOS protein levels. Furthermore, inhibition of protein farnesyltransferase with the farnesyl-protein transferase inhibitor I (0.5-50 mM) or -hydroxyfarnesylphosphonic acid (2-20 mM) did not affect ecNOS protein levels. In contrast, co-treatment with GGPP at a concentration of 10 mM, but not 1 mM, completely reversed the upregulation of ecNOS protein levels by mevastatin. These findings indicate that ecNOS expression is negatively regulated by geranylgeraniol synthesis.

#### Example 21: Effects of mevastatin on Rho membrane translocation

The geranylgeranylation of the small GTPases, RhoA and RhoB, are essential for their membrane translocation from the cytosol (Van Aelst, L and D'Souza-Schorey, C, *Genes Dev*, 1997, 11:2295-2322). Under basal culture conditions, both RhoA and RhoB are present in the membranes and cytosol. Treatment with mevastatin decreased membrane localization of RhoA and RhoB by  $60 \pm 5\%$  and  $78 \pm 6\%$  and produce a concomitant increase in RhoA and RhoB in the cytosol by  $65 \pm 4\%$  and  $87 \pm 7\%$ . Co-treatment with GGPP (5 mM), but not FPP (10 mM) reversed the effects of mevastatin and completely restored the amount of cytosolic and membrane-associated RhoA and RhoB to basal levels. These findings suggest that inhibition of Rho geranylgeranylation by mevastatin prevents RhoA and RhoB from translocating to and associating with the cellular membrane.

#### Example 22: Effects of mevastatin on Rho GTP-binding activity

To determine whether geranylgeranylation of RhoA and RhoB affects their activity (i.e. GTP-bound state), we immunoprecipitated [ $^{35}$  S]GTP S-labeled RhoA and RhoB from the membrane and cytosol of endothelial cells treated with mevastatin (10 mM) in the presence of GGPP (5 mM) or FPP (10 mM). Under basal conditions, endothelial cells have membrane-associated RhoA and RhoB activity of  $4.4 \pm 0.1$  fmol/mg/min and  $3.8 \pm 0.4$  fmol/mg/min, respectively. Treatment with mevastatin decreased membrane-associated RhoA and RhoB GTP-binding activity by 52% ( $2.1 \pm 0.4$  fmol/mg/min;  $p < 0.01$ ) and 37% ( $2.4 \pm 0.6$  fmol/mg/min;  $p < 0.05$ ), respectively ( $n = 3$ ).

Co-treatment with FPP (10 mM) produced no significant effects on RhoA and RhoB GTP-binding activity compared to mevastatin alone ( $2.6 \pm 0.9$  fmol/mg/min and  $2.7 \pm 0.5$  fmol/mg/min, respectively,  $p > 0.05$ ,  $n = 3$ ). However, co-treatment with GGPP (10 mM)



-55-

completely reversed the inhibitory effects of mevastatin on RhoA and RhoB GTP-binding activity ( $4.1 \pm 0.3$  fmol/mg/min and  $3.6 \pm 0.5$  fmol/mg/min, respectively,  $p < 0.05$ ,  $n = 3$ ). Cytosolic RhoA and RhoB were relatively inactive (i.e.  $< 1$  fmol/mg/min) and their activities were not affected by treatment with mevastatin alone or in combination with GGPP or FPP. Taken together, these results indicate that geranylgeranylation of RhoA and RhoB is necessary for their membrane translocation and that membrane-associated Rho is relatively more active in terms of GTP-binding than cytosolic Rho.

#### Example 23: Effects of C3 transferase on ecNOS expression

To determine whether the inhibition of Rho mediates the effects of mevastatin on ecNOS expression, endothelial cells were treated with mevastatin in the presence and absence of *Clostridium botulinum* C3 transferase (5-50 mg/ml), an exoenzyme which specifically inactivates Rho by ADP-ribosylation (Aktories, K, *J Clin Invest*, 1997, 12:S11-S13). Treatment of endothelial cells with mevastatin (10 mM) or C3 transferase (50 mg/ml) for 48 h augmented ecNOS protein levels by  $260 \pm 9\%$  and  $250 \pm 10\%$ , respectively ( $p < 0.01$ ,  $n = 3$ ). Lower concentrations of C3 transferase (i.e.  $< 50$  mg/ml) produced correspondingly smaller increases in ecNOS expression (data not shown). In contrast to the effect of mevastatin, the stimulatory effect of C3 transferase on ecNOS expression was not reversed in the presence of L-mevalonate (200 mM).

#### Example 24: Effects of dominant-negative RhoA on ecNOS expression

Bovine aortic endothelial cells were transfected with insertless pcDNA3 vector, c-myc-tagged wildtype RhoA (wtRhoA), or c-myc-tagged dominant-negative RhoA mutant (N19RhoA) which cannot exchange GDP for GTP and therefore is inactive (Gebbink, M et al., *J Cell Biol*, 1997, 137:1603-1613). Immunostaining for b-galactosidase activity demonstrate comparable transfection efficiency of approximately 10% among the RhoA constructs and between treatment conditions. To distinguish between transfected and endogenous RhoA, the amount of transfected RhoA constructs expressed was assessed by immunoblotting using an antibody to c-myc (9E10), which recognizes a 21 kD band only in wtRhoA and N19RhoA transfected cells (Figure 6).

Overexpression of wtRhoA mildly reduced basal ecNOS protein expression by  $15 \pm 4\%$  suggesting that increased RhoA expression results in a decrease in basal ecNOS expression ( $p < 0.05$ ,  $n = 3$ ). Endothelial cells transfected with the dominant-negative N19RhoA mutant to

comparable levels as wtRhoA as assessed by the amount of c-myc-tag, however, exhibited a 150±5% increase in ecNOS protein levels ( $p<0.05$ ,  $n=3$ ). The observed effects of N19RhoA overexpression on overall ecNOS protein levels (i.e. transfected and non-transfected cells) are more profound when one considers that only 10% of the endothelial cells were actually transfected. These findings are consistent with our earlier findings that inhibition of Rho GTPase activity leads to an increase in ecNOS expression.

Example 25: Effects of CNF-1 on ecNOS expression

The *Escherichia coli* cytotoxic necrotizing factor (CNF)-1 is known to directly and specifically activate rho proteins via glutamine deamination (Aktories, K, *J Clin Invest*, 1997, 12:S11-S13; Schmidt, G et al., *Nature*, 1997, 387:725-729; Flatau, G et al., *Nature*, 1997, 387:729-733). Treatment of endothelial cells with mevastatin (10 mM) increased ecNOS mRNA levels by  $390 \pm 15\%$  compared to basal levels ( $p<0.01$ ,  $n=3$ ). Co-treatment with CNF-1 (200 ng/ml) completely reversed the upregulation of ecNOS mRNA by mevastatin ( $p>0.05$  compared to basal levels,  $n=3$ ). Treatment with CNF-1 (200 ng/ml) alone, however, decreased ecNOS steady-state mRNA levels to  $48 \pm 6\%$  of basal levels at 24 h ( $p<0.05$ ,  $n=3$ ). These findings indicate that the direct activation of Rho leads to the downregulation of ecNOS expression.

Example 26: Effects of HMG-CoA Reductase Inhibitors and Rho on ecNOS activity

The ecNOS activity was assessed by measuring the LNMA-inhibitable nitrite accumulation in conditioned media of endothelial cells (Laufs, U et al., *J Biol Chem*, 1997, 272:31725-31729). Basal ecNOS activity was  $9.7 \pm 1.4$  nmol/500,000 cells/24 hours (Figure 7). Treatment of endothelial cells with mevastatin (10 mM) resulted in a 3-fold increase in nitrite accumulation ( $32 \pm 1.9$  nmol/500,000 cells/24 hours,  $p<0.01$ ). This increase in ecNOS activity by mevastatin was reversed by co-treatment with GGPP (5 mM), but not FPP (10 mM) ( $12 \pm 0.8$  and  $27 \pm 4.9$  nmol/500,000 cells/24 hours, respectively). Furthermore, direct activation of Rho by CNF-1 (200 ng/ml) reversed mevastatin-induced increase in ecNOS activity ( $32 \pm 1.9$  to  $14 \pm 2.1$  nmol/500,000 cells/24 hours,  $p<0.05$ ). In contrast, inhibition of Rho by C3 transferase (50 mg/ml) resulted in a 3-fold increase in nitrite accumulation ( $31 \pm 2.1$  and  $\pm$  nmol/500,000 cells/24 hours,  $p<0.05$ ). These results indicate that Rho not only negatively regulates ecNOS expression, but also ecNOS activity.

*"Regulation of Endothelial Nitric Oxide Synthase Activity by Agents that Disrupt Actin Cytoskeletal Organization"*

Experimental Procedures

5 Materials:

Myosin light chain kinase inhibitors BDM [2,3-butanedione 2-monoxime], ML-7 [1-(5-iodonaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride], and H-7 [1-(5-isoquinolinesulphonyl)-2-methylpiperazine dihydro-chloride], were purchased from Sigma Chemical Corp. (St. Louis, MO). Nocodazole {Methyl-(5-[2-thienylcarbonyl]-1H-benzimidazol-2-yl)carbamate} was also purchased from Sigma.

Cell culture. Western blotting and Northern blotting were performed as described earlier.

Example 27: Effects of Agents That Disrupt Actin Cytoskeletal Organization on ecNOS Protein and mRNA Expression

15 To determine whether downstream targets of *rho*GTPase exert any effect on ecNOS expression, endothelial cells were treated in the presence and absence of a myosin light chain (MLC) kinase inhibitor, for example, H-7. MLC kinase inhibitors decrease MLC phosphorylation and stress fiber formation. Treatment of endothelial cells with H-7 (1-100 mM) for 24 hours augmented ecNOS protein levels (Figure 8). Similar experiments using a different  
20 MLC kinase inhibitor (ML-7), produced identical results.

Furthermore, disruption of the actin cytoskeletal organization by Cytochalasin D (an agent that interferes with actin polymerization) or BDM (a different MLC kinase inhibitor), also lead to upregulation of ecNOS (Figures 9 and 10 respectively).

25 Inhibition, however, of cell cycle progression by anaphicoline (data not shown) or disruption of the microtubular cytoskeleton by enhancing microtubule depolymerization with nocodazole, do not increase ecNOS expression (Figure 11).

These findings show that the upregulation of ecNOS expression is relatively specific to disruption of the actin cytoskeleton. Additionally, the increased ecNOS expression by Cytochalasin D or BDM does not result from nonspecific cytotoxicity of these agents.

30 Example 28: Effect of L-arginine on cerebral blood flow

L-arginine infusion at 300 mg/kg, i.v., caused modest (10%) and variable elevations in

regional cerebral blood flow (rCBF) after infusion in several preliminary experiments (n=4, data not shown). In the present experiments, 450 mg/kg or saline was infused at a constant rate of 100 microliter/kg/min over 15 minutes into wild type mice, mutant mice deficient in endothelial nitric oxide synthase (eNOS null), and mice which had received chronic daily administration of simvastatin (2 mg/kg). Regional cerebral blood flow (rCBF) was monitored by laser-Doppler flowimetry in groups of urethane-anesthetized, ventilated mice. Additional physiological variables were also monitored in the mice, including mean arterial blood pressure (MABP), heart rate, blood pH, PaO<sub>2</sub>, and PaCO<sub>2</sub>.

### Results

Physiological variables during laser-Doppler flowimetry in urethane-anesthetized ventilated wild type, simvastatin-treated and eNOS null mice infused with L-arginine or saline are shown in Table 1. Number of mice in each group is shown in parenthesis. Values are reported as mean +/- SEM. \* denotes statistically significant difference (P<0.05) compared with eNOS null mice; # denotes statistically significant difference (P<0.05) compared with baseline by one-way ANOVA followed by Scheffe test. MABP indicates mean arterial blood pressure; sim indicates mice chronically administered simvastatin.

There were no within-group differences during observation time in mean arterial blood pressure and heart rate, although those values were elevated in eNOS null mice as reported previously. PaCO<sub>2</sub> values were not different between two time points in all groups nor between-group, although pH values were lower after infusion of L-arginine.

#### *rCBF response to L-arginine:*

Figure 21 is a bar graph showing regional CBF changes in wild type and eNOS null mice for 40 min after L-arginine (450 mg/kg) or saline infusion at a constant rate of 100 microliter/kg/min over 15 min. The number of mice in each group is indicated in parenthesis. Error bars denote standard error of the mean (SEM), and an asterisk (\*) denotes statistically significant difference (P<0.05) compared with baseline control by one-way ANOVA followed by Fisher's protected least-squares difference test.

L-arginine infusion (450 mg/kg, i.v.) increased rCBF in parietal cortex in wild type mice, as shown in Figure 1 (Fig. 1). The increase in rCBF began at 5-10 minutes and achieved statistical significance at 10-15 minutes after infusion. Maximum values achieved at 20-25 min reached 26% above, after which values decreased to control levels. By contrast, L-arginine did not increase rCBF in eNOS null mice. Values in these mutants ranged from -4 to +5% during the

40 minute recording period. Saline infusion in wild type mice did not increase rCBF significantly.

*rCBF response to L-arginine plus simvastatin:*

Figure 22 is a bar graph showing regional CBF changes in simvastatin-treated mice for 40 min after L-arginine or saline infusion at the same dose. The number of mice in each group is indicated in parenthesis; sim indicates simvastatin. Error bars denote SEM and an asterisk (\*) denotes statistically significant difference ( $P < 0.05$ ) compared with baseline control by one-way ANOVA followed by Fisher's protected least-squares difference test.

After chronic daily administration of simvastatin alone, the baseline rCBF was increased by 25%. L-arginine but not saline infusions increased rCBF significantly above the simvastatin baseline. Marked elevation was observed in the 10-15 minute epoch. The maximum increase was observed at 15-20 min and was 29-31% over baseline. These increases sustained for an additional 20 minutes which was considerably longer than after L-arginine treatment alone. The maximum response to L-arginine in the presence of simvastatin was not statistically increased. However, the response to L-arginine was more sustained in the simvastatin-treated mice. In the 30-40 minute epoch, the increase in blood flow was larger in the simvastatin treated compared to non treated control ( $P < 0.05$ ).

TABLE 1

	MABP	heart rate	pH	PaO <sub>2</sub>	PaCO <sub>2</sub>
Group (n)	mmHg	bpm		mmHg	mmHg
wild + saline (6)					
baseline	94.7 $\pm$ 3.4	543 $\pm$ 20	7.36 $\pm$ 0.02	154 $\pm$ 13	35.8 $\pm$ 2.0
0-5 min	94.8 $\pm$ 3.3	549 $\pm$ 19			
10-15 min	96.2 $\pm$ 3.2	548 $\pm$ 16			
20-25 min	95.8 $\pm$ 3.1	550 $\pm$ 16			
35-40 min	96.5 $\pm$ 2.9	547 $\pm$ 15			
after infusion			7.35 $\pm$ 0.02	180 $\pm$ 5	33.4 $\pm$ 1.8
wild + L-arginine (7)					
baseline	92.9 $\pm$ 3.5*	545 $\pm$ 11	7.40 $\pm$ 0.02	127 $\pm$ 9	39.2 $\pm$ 2.0
0-5 min	92.7 $\pm$ 3.5	548 $\pm$ 11			
10-15 min	94.6 $\pm$ 3.5	561 $\pm$ 9			

-60-

	20-25 min	93.3+/-3.4	554+/-10			
	35-40 min	89.9+/-3.3	533+/-9			
	after infusion			7.32+/-0.03	169+/-5#	36.4+/-1.7
	eNOS null+L-arginine (4)					
5	baseline	116.3+/-9.7	618+/-9	7.40+/-0.03	15+/-4	36.4+/-1.7
	0-5 min	115.8+/-10.0	621+/-8			
	10-15 min	113.3+/-7.4	623+/-6			
	20-25 min	113.8+/-8.1	630+/-13			
	35-40 min	94.8+/-8.1	604+/-10			
10	after infusion			7.28+/-0.04#	178+/-7#	35.8+/-2.4

-61-

Table 1 continued—					
sim (2mg/kg)+saline (3)					
baseline	88.0+/-3.0*	541+/-1	7.46+/-0.03	159+/-19	32.7+/-2.2
0-5 min	90.7+/-2.8	543+/-3			
10-15 min	93.3+/-3.3	547+/-9			
20-25 min	95.0+/-3.5	553+/-10			
35-40 min	94.0+/-4.0	558+/-4			
after infusion			7.41+/-0.01	177+/-7	32.5+/-2.8
sim (2mg/kg)+L-arginine(5)					
baseline	87.4+/-3.1*	505+/-9*	7.44+/-0.03	144+/-15	31.2+/-2.8
0-5 min	88.4+/-3.3*	503+/-7*			
10-15 min	92.2+/-3.2	507+/-6*			
20-25 min	88.4+/-3.4*	502+/-5*			
35-40 min	83.8+/-4.2	490+/-4*			
after infusion			7.30+/-0.02#	163+/-13	34.2+/-2.1
sim (20mg/kg)+L-arginine (6)					
baseline	91.7+/-2.8*	566+/-26	7.44+/-0.01	169+/-8	32.0+/-1.5
0-5 min	91.5+/-3.7*	571+/-26			
10-15 min	92.7+/-4.9	574+/-26			
20-25 min	89.7+/-4.9*	571+/-26			
35-40 min	84.8+/-5.0	551+/-21			
after infusion			7.33+/-0.02#	178+/-7	32.4+/-2.7

### Detailed description of the drawings

Figure 1. A) NOS-Activity measured by (C14)-arginine-citrulline assay in the aortas of wild-type SV-129 mice after treatment with simvastatin (Sim, 2 mg/kg, s.c., 14 days) and of mice injected with PBS (Control), n = 8, p < 0.05.

B) ecNOS mRNA expression determined by quantitative polymerase chain reaction in wild-type SV-129 mice aortas after treatment with simvastatin (Sim0.2, 0.2 mg/kg, s.c. and Sim20, 20 mg/kg, s.c.) for 14 days and of mice injected with saline (Control) in comparison to glyceraldehyde 3-phosphate dehydrogenase (G3DPH) mRNA expression.

ecNOS expression and function is upregulated in the aortas of mice treated with Sim.

**Figure 2.** ecNOS mRNA expression in the infarcted, ipsilateral (I) and not-infarcted, contralateral (C) forebrain hemispheres of SV-129 mice after treatment with simvastatin (Sim, 2 mg/kg, s.c., 14 days) and mice injected with saline (Control), as determined by quantitative polymerase chain reaction compared to glyceraldehyde 3-phosphate dehydrogenase (G3DPH) mRNA expression. ecNOS mRNA expression was upregulated in the infarcted brain area in Sim-treated animals.

**Figure 3. A)** Northern analyses (20 mg total RNA/lane) showing the effects of mevastatin (Statin, 10 mM) alone or in combination with FPP (10 mM) or GGPP (10 mM) on eNOS steady-state mRNA levels at 24 h. **B)** Concentration-dependent effects of GGPP (1-10 mM) on mevastatin (10 mM)-induced increases in eNOS mRNA levels after 24 h. Each experiment was performed three times with comparable results. The corresponding ethidium bromide-stained 28S band intensities were used to standardize loading conditions.

**Figure 4.** Immunoblots (30 mg protein/lane) showing the effects of mevastatin (Statin, 10 mM) alone or in combination with FPP (10 mM), GGPP (1 - 10 mM), or LDL cholesterol (LDL-C, 1 mg/ml) on eNOS protein levels after 24 h. The blot is representative of three separate experiments.

**Figure 5.** Immunoblot (30 mg protein/lane) showing the effects of C3 transferase (C3, 50 mg/ml), mevastatin (Statin, 10 mM), or L-mevalonate (Mev, 200 mM) on eNOS protein levels after 48 h. The blot is representative of three separate experiments.

**Figure 6.** Immunoblots (30 mg protein/lane) showing eNOS protein levels after transfection with insertless vector, pcDNA3 (C), c-myc-wildtype-RhoA (wt), and c-myc-N19RhoA (dominant-negative rhoA mutant). The levels of overexpressed RhoA mutants were determined by immunoblotting for their corresponding c-myc-tags (c-myc-RhoA). Experiments were performed three times with similar results.

**Figure 7.** Effects of C3 transferase (C3, 50 mg/ml), FPP (10 mM), GGPP (5 mM), and CNF-1 (200 ng/ml) on mevastatin (Statin, 10 mM)-induced eNOS activity as determined by LNMA-inhibitable nitrite production at 24 h. Experiments were performed three times in duplicate



-63-

with less than 5% variation. \* $p < 0.05$  compared with control (C), \*\* $p < 0.05$  compared with mevastatin.

5 Figure 8. Immunoblots (30 mg protein/lane) showing the concentration-dependent effects of MLC kinase inhibitor H-7 on eNOS protein levels after 24 hours.

Figure 9. Northern blot analysis (20 mg total RNA/lane) showing eNOS mRNA expression of endothelial cells treated with cytochalasin D at 24 hours.

10 Figure 10. Immunoblots (30 mg protein/lane) showing the concentration-dependent effects of 2, 3-butanedione 2-monoxime on eNOS protein levels.

Figure 11. Northern blot analysis (20 mg total RNA/lane) showing eNOS mRNA expression of endothelial cells treated with nocodazole for 24 hours.

15 Figure 12. Bar graph showing regional CBF changes in wild type and eNOS null mice for 40 min after L-arginine or saline infusion.

20 Figure 13. Bar graph showing regional CBF changes in simvastatin-treated mice for 40 min after L-arginine or saline infusion at the same dose.

A Sequence Listing follows the claims. All references disclosed herein are incorporated by reference in their entirety.

I claim:

25

-64-

Claims

1. A method for increasing endothelial cell Nitric Oxide Synthase activity in a subject who would benefit from increased endothelial cell Nitric Oxide Synthase activity in a tissue comprising:

5 administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

10 2. The method of claim 1, wherein the agent that disrupts actin cytoskeletal organization is not H-7, or fasudil, when the subject in need of such treatment has an abnormally elevated risk of an ischemic stroke

15 3. The method of claim 1, wherein the subject is nonhypercholesterolemic.

4 The method of claim 1, wherein the subject is nonhyperlipidemic.

5. The method of claim 1, wherein the amount is sufficient to increase endothelial cell Nitric Oxide Synthase activity above normal baseline levels.

20 6. The method of claim 1, wherein the subject has a condition comprising an abnormally low level of endothelial cell Nitric Oxide Synthase activity which is chemically induced.

25 7. The method of claim 1, wherein the subject has an abnormally elevated risk of pulmonary hypertension.

8. The method of claim 1, wherein the subject has pulmonary hypertension.

30 9. The method of claim 1, wherein the subject has an abnormally elevated risk of an ischemic stroke.

10. The method of claim 1, wherein the subject has experienced an ischemic stroke.

-65-

11. The method of claim 1, wherein the subject is chronically exposed to hypoxic conditions.
12. The method of claim 1, wherein the subject has an abnormally elevated risk of thrombosis.
- 5 13. The method of claim 1, wherein the subject has thrombosis.
14. The method of claim 1, wherein the subject has an abnormally elevated risk of arteriosclerosis.
- 10 15. The method of claim 1, wherein the subject has arteriosclerosis.
16. The method of claim 1, wherein the subject has an abnormally elevated risk of myocardial infarction.
- 15 17. The method of claim 1, wherein the subject has experienced a myocardial infarction.
18. The method of claim 1, wherein the subject has an abnormally elevated risk of reperfusion injury.
- 20 19. The method of claim 18, wherein the subject is a transplant recipient.
20. The method of claim 1, wherein the subject has homocystinuria.
21. The method of claim 1, wherein the subject has a neurodegenerative disease.
- 25 22. The method of claim 21, wherein the neurodegenerative disease is Alzheimer's disease.
23. The method of claim 1, wherein the subject has CADASIL syndrome.
- 30 24. The method of claim 1, wherein the agent that disrupts actin cytoskeletal organization is not H-7, when the subject in need of such treatment has an abnormally elevated risk of an

-66-

ischemic stroke.

25. The method of claim 1, wherein the agent that disrupts actin cytoskeletal organization is not fasudil, when the subject in need of such treatment has an abnormally elevated risk of an ischemic stroke.

26. The method of claims 1- 25, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

27. The method of claim 26, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

28. The method of claim 26, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

29. The method of claim 28, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

30. The method of claims 1-25, wherein the agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D.

31. The method of claim 30, wherein the myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)-2-methylpiperazine dihydro-chloride.

32. The method of claim 30, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

33. The method of claim 32, wherein the endothelial cell Nitric Oxide Synthase substrate is

-67-

L-arginine.

34. The method of claim 32, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

5

35. The method of claim 34, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

10

36. The method of claims 1-25, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject.

15

37. The method of claim 36, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

38. The method of claim 37, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

20

39. The method of claim 37, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

40. The method of claim 39, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

25

41. The method of claims 1-25, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization that increases endothelial cell Nitric Oxide Synthase activity.

30

42. The method of claim 41, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

43. The method of claim 42, wherein the endothelial cell Nitric Oxide Synthase substrate is

-68-

L-arginine.

44. The method of claim 42, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

5

45. The method of claim 44, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

10

46. A method for increasing endothelial cell Nitric Oxide Synthase activity in a subject to treat a condition favorably affected by an increase in endothelial cell Nitric Oxide Synthase activity in a tissue comprising:

15

administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

20

47. The method of claim 46, wherein the agent that disrupts actin cytoskeletal organization is not H-7, or fasudil, when the subject in need of such treatment has an abnormally elevated risk of an ischemic stroke.

48. The method of claim 46, wherein the subject is nonhypercholesterolemic.

49. The method of claim 46, wherein the subject is nonhyperlipidemic.

25

50. The method of claim 46, wherein the amount is sufficient to increase endothelial cell Nitric Oxide Synthase activity above normal baseline levels.

51. The method of claim 46, wherein the subject has a condition comprising an abnormally low level of endothelial cell Nitric Oxide Synthase activity which is chemically induced.

30

52. The method of claim 46, wherein the subject has an abnormally elevated risk of pulmonary hypertension.

-69-

53. The method of claim 46, wherein the subject has pulmonary hypertension.

54. The method of claim 46, wherein the subject has an abnormally elevated risk of an ischemic stroke.

5

55. The method of claim 46, wherein the subject has experienced an ischemic stroke.

56. The method of claim 46, wherein the subject is chronically exposed to hypoxic conditions.

10

57. The method of claim 46, wherein the subject has an abnormally elevated risk of thrombosis.

58. The method of claim 46, wherein the subject has thrombosis.

15

59. The method of claim 46, wherein the subject has an abnormally elevated risk of arteriosclerosis.

60. The method of claim 46, wherein the subject has arteriosclerosis.

20

61. The method of claim 46, wherein the subject has an abnormally elevated risk of myocardial infarction.

62. The method of claim 46, wherein the subject has experienced a myocardial infarction.

25

63. The method of claim 46, wherein the subject has an abnormally elevated risk of reperfusion injury.

64. The method of claim 63, wherein the subject is a transplant recipient.

30

65. The method of claim 46, wherein the subject has homocystinuria.

66. The method of claim 46, wherein the subject has a neurodegenerative disease.

-70-

67. The method of claim 66, wherein the neurodegenerative disease is Alzheimer's disease.

68. The method of claim 46, wherein the subject has CADASIL syndrome.

69. The method of claim 46, wherein the agent that disrupts actin cytoskeletal organization is not H-7, when the subject in need of such treatment has an abnormally elevated risk of an ischemic stroke.

70. The method of claim 46, wherein the agent that disrupts actin cytoskeletal organization is not fasudil, when the subject in need of such treatment has an abnormally elevated risk of an ischemic stroke.

71. The method of claims 46-70, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

72. The method of claim 71, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

73. The method of claim 71, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

74. The method of claim 73, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

75. The method of claims 46-70, wherein the agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D.

76. The method of claim 75, wherein the myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-



-71-

1H-hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)-2-methylpiperazine dihydro-chloride.

5 77. The method of claim 75, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

78. The method of claim 77, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

10 79. The method of claim 77, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

80. The method of claim 79, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

15 81. The method of claims 46-70, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject.

20 82. The method of claim 81, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

83. The method of claim 82, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

25 84. The method of claim 82, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

30 85. The method of claim 84, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

86. The method of claims 46-70, further comprising co-administering a non-agent that

-72-

disrupts actin cytoskeletal organization agent that increases endothelial cell Nitric Oxide Synthase activity.

87. The method of claim 86, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

88. The method of claim 87, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

89. The method of claim 87, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

90. The method of claim 89, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

91. A method for reducing brain injury resulting from a stroke, comprising:

administering to a subject having an abnormally high risk of an ischemic stroke, an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the brain tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

92. The method of claim 91, wherein the agent that disrupts actin cytoskeletal organization is not H-7, or fasudil when the subject in need of such treatment has an abnormally elevated risk of an ischemic stroke.

93. The method of claim 91, wherein the subject is nonhypercholesterolemic.

94. The method of claim 91, wherein the subject is nonhyperlipidemic.

95. The method of claim 91, wherein the agent that disrupts actin cytoskeletal organization is administered prophylactically.

-73-

96. The method of claim 91, wherein the agent that disrupts actin cytoskeletal organization is administered acutely.

97. The method of claims 91-96, wherein the agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D.

98. The method of claim 97, wherein the myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)-2-methylpiperazine dihydro-chloride.

99. The method of claims 91-96, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

100. The method of claim 99, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

101. The method of claim 99, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

102. The method of claim 101, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

103. The method of claims 91-96, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization that increases endothelial cell Nitric Oxide Synthase activity.

104. The method of claim 103, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

-74-

105. The method of claim 104, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

106. The method of claim 104, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

107. The method of claim 106, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

108. The method of claims 91-96, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject.

109. The method of claim 108, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

110. The method of claim 109, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

111. The method of claim 109, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

112. The method of claim 111, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

113. A method for treating pulmonary hypertension comprising:

administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the pulmonary tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

114. The method of claim 113, wherein the subject is nonhypercholesterolemic.

-75-

115. The method of claim 113, wherein the subject is nonhyperlipidemic.

116. The method of claim 113, wherein the *rho* GTPase function inhibitor is administered prophylactically to a subject who has an abnormally elevated risk of developing pulmonary hypertension.

117. The method of claim 113, wherein the agent that disrupts actin cytoskeletal organization is administered acutely to a subject who has pulmonary hypertension.

118. The method of claim 113, wherein the subject is chronically exposed to hypoxic conditions.

119. The method of claims 113-118, wherein the agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D.

120. The method of claim 119, wherein the myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)-2-methylpiperazine dihydro-chloride.

121. The method of claims 113-118, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

122. The method of claim 121, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

123. The method of claim 121, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

124. The method of claim 123, wherein the endothelial cell Nitric Oxide Synthase cofactor is

-76-

NADPH or tetrahydrobiopterin.

125. The method of claims 113-118, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization that increases endothelial cell Nitric Oxide Synthase activity.

126. The method of claims 113-118, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject.

127. The method of claim 125, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

128. The method of claim 127, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

129. The method of claim 127, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

130. The method of claim 129, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

131. A method for treating heart failure comprising:  
administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the heart tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

132. The method of claim 131, wherein the subject is nonhypercholesterolemic.

133. The method of claim 131, wherein the subject is nonhyperlipidemic.

-77-

134. The method of claim 131, wherein the agent that disrupts actin cytoskeletal organization is administered prophylactically to a subject who has an abnormally elevated risk of heart failure.

135. The method of claim 131, wherein the agent that disrupts actin cytoskeletal organization is administered acutely to a subject who has heart failure.

136. The method of claims 131-135, wherein the agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D.

137. The method of claim 136, wherein the myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)-2-methylpiperazine dihydro-chloride.

138. The method of claims 136, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

139. The method of claim 138, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

140. The method of claim 138, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

141. The method of claim 140, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

142. The method of claims 131-135, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject.

-78-

143. The method of claim 142, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

5 144. The method of claim 143, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

145. The method of claim 143, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

10 146. The method of claim 145, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

147. The method of claims 131-135, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

15 148. The method of claim 147, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

20 149. The method of claim 147, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

150. The method of claim 149, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

25 151. The method of claims 131-135, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization that increases endothelial cell Nitric Oxide Synthase activity.

30 152. The method of claim 151, further comprising co-administering an endothelial cell Nitric Oxide Synthase substrate.

153. The method of claim 152, wherein the endothelial cell Nitric Oxide Synthase substrate



-79-

is L-arginine.

154. The method of claim 152, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

5

155. The method of claim 154, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

156. A method for treating progressive renal disease comprising:

10

administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the kidney tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

15

157. The method of claim 156, wherein the subject is nonhypercholesterolemic.

158. The method of claim 156, wherein the subject is nonhyperlipidemic.

20

159. The method of claim 156, wherein the agent that disrupts actin cytoskeletal organization is administered prophylactically.

160. The method of claim 156, wherein the agent that disrupts actin cytoskeletal organization is administered acutely.

25

161. The method of claims 156-160, wherein the agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D.

30

162. The method of claim 161, wherein the myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)-2-methylpiperazine

-80-

dihydro-chloride.

163. The method of claims 156-160, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

5

164. The method of claims 156-160, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization that increases endothelial cell Nitric Oxide Synthase activity.

10

165. The method of claims 156-160, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject.

15

166. The method of claim 164, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

20

167. A method for increasing blood flow in a tissue of a subject, comprising administering to a subject in need of such treatment an agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in the tissue of the subject, provided that the agent that disrupts actin cytoskeletal organization is not a *rho* GTPase function inhibitor.

25

168. The method of claim 167, wherein blood flow is increased in brain tissue.

169. The method of claim 167 wherein the subject is nonhypercholesterolemic.

170. The method of claim 167, wherein the subject is nonhyperlipidemic.

171. The method of claim 168, wherein the subject is nonhypercholesterolemic.

30

172. The method of claim 168, wherein the subject is nonhyperlipidemic.

-81-

173. The method of claim 167, wherein the agent that disrupts actin cytoskeletal organization is administered prophylactically.

174. The method of claim 167, wherein the agent that disrupts actin cytoskeletal organization is administered acutely.

175. The method of claims 167-174, wherein the agent that disrupts actin cytoskeletal organization is selected from the group consisting of a myosin light chain kinase inhibitor, a myosin light chain phosphatase, a protein kinase N inhibitor, a phosphatidylinositol 4-phosphate 5-kinase inhibitor, and cytochalasin D.

176. The method of claim 175, wherein the myosin light chain kinase inhibitor is selected from the group consisting of 2,3-butanedione 2-monoxime, 1-(5-iodonaphthalene-1-sulphonyl)-1H-hexahydro-1,4-diazepine hydrochloride, and 1-(5-isoquinolinesulphonyl)-2-methylpiperazine dihydro-chloride.

177. The method of claims 167-174, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

178. The method of claim 177 wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

179. The method of claim 177, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

180. The method of claim 179, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

181. The method of claims 167-174, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization that increases endothelial cell Nitric Oxide Synthase activity.

-82-

182. The method of claims 167-174, further comprising co-administering at least one different agent that disrupts actin cytoskeletal organization in an amount effective to increase endothelial cell Nitric Oxide Synthase activity in said tissue of the subject.

5 183. The method of claim 181, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

184. The method of claim 183, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

10 185. The method of claim 183, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

15 186. The method of claim 185, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

20 187. The method of claims 167-174, further comprising co-administering a second agent to the subject with a condition treatable by the second agent in an amount effective to treat the condition, whereby the delivery of the second agent to a tissue of the subject is enhanced as a result of the increased blood flow.

188. The method of claim 187, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

25 189. The method of claim 188, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

190. The method of claim 188, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

30 191. The method of claim 190, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

-83-

192. The method of claims 167-174, further comprising co-administering a second agent to the subject with a condition treatable by the second agent in an amount effective to treat the condition, whereby the delivery of the second agent to the brain of the subject is enhanced as a result of the increased blood flow.

5

193. The method of claims 167-174, further wherein the tissue is brain and the second agent comprises an agent having a site of action in the brain.

10

194. The method of claim 192, wherein the second agent is selected from the group consisting of analeptic, analgetic, anesthetic, adrenergic agent, anti-adrenergic agent, amino acids, antagonists, antidote, anti-anxiety agent, anticholinergic, anticolvunsant, antidepressant, anti-emetic, anti-epileptic, antihypertensive, antifibrinolytic, antihyperlipidemia, antimigraine, antinauseant, antineoplastic (brain cancer), antiobessional agent, antiparkinsonian, antipsychotic, appetite suppressant, blood glucose regulator, cognition adjuvant, cognition enhancer, dopaminergic agent, emetic, free oxygen radical scavenger, glucocorticoid, hypocholesterolemic, hyperlipidemic, histamine H2 receptor antagonists, immunosuppressant, inhibitor, memory adjuvant, mental performance enhancer, mood regulator, mydriatic, neuromuscular blocking agent, neuroprotective, NMDA antagonist, post-stroke and post-head trauma treatment, psychotropic, sedative, sedative-hypnotic, serotonin inhibitor, tranquilizer, and treatment of cerebral ischemia, calcium channel blockers, free radical scavengers - antioxidants, GABA agonists, glutamate antagonists, AMPA antagonists, kainate antagonists, competitive and non-competitive NMDA antagonists, growth factors, opioid antagonists, phosphatidylcholine precursors, serotonin agonists, sodium- and calcium-channel blockers, and potassium channel openers.

15

20

25

195. The method of claim 193, wherein the agent having a site of action in the brain is selected from the group consisting of analeptic, analgetic, anesthetic, adrenergic agent, anti-adrenergic agent, amino acids, antagonists, antidote, anti-anxiety agent, anticholinergic, anticolvunsant, antidepressant, anti-emetic, anti-epileptic, antihypertensive, antifibrinolytic, antihyperlipidemia, antimigraine, antinauseant, antineoplastic (brain cancer), antiobessional agent, antiparkinsonian, antipsychotic, appetite suppressant, blood glucose regulator, cognition adjuvant, cognition enhancer, dopaminergic agent, emetic, free oxygen radical scavenger, glucocorticoid,

30

-84-

hypocholesterolemic, hollylipidemic, histamine H2 receptor antagonists, immunosuppressant, inhibitor, memory adjuvant, mental performance enhancer, mood regulator, mydriatic, neuromuscular blocking agent, neuroprotective. NMDA antagonist, post-stroke and post-head trauma treatment, psychotropic, sedative, sedative-hypnotic, serotonin inhibitor, tranquilizer, and treatment of cerebral ischemia, calcium channel blockers, free radical scavengers - antioxidants, GABA agonists, glutamate antagonists, AMPA antagonists, kainate antagonists, competitive and non-competitive NMDA antagonists, growth factors, opioid antagonists, phosphatidylcholine precursors, serotonin agonists, sodium- and calcium-channel blockers, and potassium channel openers.

196. The method of claim 187, wherein the second agent is selected from the group consisting of analeptic, analgetic, anesthetic, adrenergic agent, anti-adrenergic agent, amino acids, antagonists, antidote, anti-anxiety agent, anticholinergic, anticolvunsant, antidepressant, anti-emetic, anti-epileptic, antihypertensive, antifibrinolytic, antihyperlipidemia, antimigraine, antinauseant, antineoplastic (brain cancer), antiobessional agent, antiparkinsonian, antipsychotic, appetite suppressant, blood glucose regulator, cognition adjuvant, cognition enhancer, dopaminergic agent, emetic, free oxygen radical scavenger, glucocorticoid, hypocholesterolemic, hollylipidemic, histamine H2 receptor antagonists, immunosuppressant, inhibitor, memory adjuvant, mental performance enhancer, mood regulator, mydriatic, neuromuscular blocking agent, neuroprotective, NMDA antagonist, post-stroke and post-head trauma treatment, psychotropic, sedative, sedative-hypnotic, serotonin inhibitor, tranquilizer, and treatment of cerebral ischemia, calcium channel blockers, free radical scavengers - antioxidants, GABA agonists, glutamate antagonists, AMPA antagonists, kainate antagonists, competitive and non-competitive NMDA antagonists, growth factors, opioid antagonists, phosphatidylcholine precursors, serotonin agonists, sodium- and calcium-channel blockers, and potassium channel openers.

197. The method of claim 192, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

198. The method of claim 197, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

-85-

199. The method of claim 197, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

200. The method of claim 199, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

201. The method of claim 193, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

202. The method of claim 201, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

203. The method of claim 201, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

204. The method of claim 203, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.

205. A method of screening for identifying an agent that disrupts actin cytoskeletal organization for treatment of subjects who would benefit from increased endothelial cell Nitric Oxide Synthase activity in a tissue, comprising:

(a) identifying an agent that disrupts actin cytoskeletal organization suspected of increasing endothelial cell Nitric Oxide Synthase activity, and

(b) determining whether or not the agent that disrupts actin cytoskeletal organization produces an increase in endothelial cell Nitric Oxide Synthase activity *in vivo* or *in vitro*.

206. The method of claim 205, wherein the subject who would benefit from increased endothelial cell Nitric Oxide Synthase activity in a tissue has an abnormally elevated risk of an ischemic stroke or has experienced a stroke, and the increased endothelial cell Nitric Oxide Synthase activity is increased in brain tissue.

-86-

207. A composition comprising an agent that disrupts actin cytoskeletal organization and L-arginine.

5 208. The composition according to claim 207, wherein the composition is a pharmaceutical composition.

209. The composition according to claim 207, wherein the agent that disrupts actin cytoskeletal organization and the L-arginine are in amounts effective to increase blood flow.

10 210. The composition according to claim 207, wherein the agent that disrupts actin cytoskeletal organization and the L-arginine are in amounts effective to increase blood flow in brain tissue.

211. The composition according to claim 207, wherein the administration of said composition results in increased blood flow.

15 212. The composition according to claim 207, wherein the administration of said composition results in increased blood flow to the brain.

20 213. The method of claim 182, further comprising co-administering a substrate of endothelial cell Nitric Oxide Synthase.

214. The method of claim 213, wherein the endothelial cell Nitric Oxide Synthase substrate is L-arginine.

25 215. The method of claim 213, further comprising co-administering an endothelial cell Nitric Oxide Synthase cofactor.

30 216. The method of claim 215, wherein the endothelial cell Nitric Oxide Synthase cofactor is NADPH or tetrahydrobiopterin.



1/8




Fig. 1A

## Aortic ecNOS mRNA Expression




Fig. 1B

SUBSTITUTE SHEET (RULE 26)





Fig. 2

3/8



18 S —				
Statin	-	+	+	+
FPP	-	-	+	-
GGPP	-	-	-	+

Fig. 3A




Fig. 3B




Fig. 4

SUBSTITUTE SHEET (RULE 26)

4/8




Fig. 5




Fig. 6




Fig. 7

SUBSTITUTE SHEET (RULE 26)

5/8




Fig. 8




Fig. 9

6/8




Fig. 10




Fig. 11

7/8




Fig. 12

8/8




Fig. 13



## SEQUENCE LISTING

<110> The Brigham and Women's Hospital, Inc.

<120> UPREGULATION OF TYPE III ENDOTHELIAL  
CELL NITRIC OXIDE SYNTHASE BY AGENTS THAT DISRUPT ACTIN  
CYTOSKELETAL ORGANIZATION

<130> R0547/7008WO/ERG/KA

<150> US 09/115,387

<151> 1998-07-14

<150> US 09/273,224

<151> 1999-03-19

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 25

<212> DNA

<213> Mus Musculus

<400> 1

gggctccctc cttccggctg ccacc

25

<210> 2

<211> 24

<212> DNA

<213> Mus Musculus

<400> 2

ggatccctgg aaaaggcggt gagg

24

<210> 3

<211> 20

<212> DNA

<213> Mus Musculus

<400> 3

accacagtcc atgccatcac

20

<210> 4

<211> 20

<212> DNA

<213> Mus Musculus

<400> 4

tccaccaccc tgttgctgta

20




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 31/15, 31/4035, 31/496, 31/551, A61P 9/10, 7/00</b>		<b>A3</b>	(11) International Publication Number: <b>WO 00/03746</b>
			(43) International Publication Date: 27 January 2000 (27.01.00)
(21) International Application Number: PCT/US99/15827		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 14 July 1999 (14.07.99)		Published With international search report.	
(30) Priority Data: 09/115,387 14 July 1998 (14.07.98) US 09/273,224 19 March 1999 (19.03.99) US		(88) Date of publication of the international search report: 20 April 2000 (20.04.00)	
(71) Applicant: THE BRIGHAM AND WOMEN'S HOSPITAL, INC. [US/US]; 75 Francis Street, Boston, MA 02115 (US).			
(72) Inventor: LIAO, James, K.; 12 Audubon Road, Weston, MA 02193 (US).			
(74) Agent: GATES, Edward, R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).			

(54) Title: UPREGULATION OF TYPE III ENDOTHELIAL CELL NITRIC OXIDE SYNTHASE BY AGENTS THAT DISRUPT ACTIN CYTOSKELETAL ORGANIZATION

## (57) Abstract

A use for agents that disrupt actin cytoskeletal organization is provided. In the instant invention, agents that disrupt actin cytoskeletal organization are found to upregulate endothelial cell Nitric Oxide Synthase activity. As a result, agents that disrupt actin cytoskeletal organization are useful in treating or preventing conditions that result from the abnormally low expression and/or activity of endothelial cell Nitric Oxide Synthase. Such conditions include pulmonary hypertension, ischemic stroke, impotence, heart failure, hypoxia-induced conditions, insulin deficiency, progressive renal disease, gastric or esophageal motility syndrome, etc. Subjects thought to benefit mostly from such treatments include nonhyperlipidemics and nonhypercholesterolemics, but not necessarily exclude hyperlipidemics and hypercholesterolemics.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# INTERNATIONAL SEARCH REPORT

Intern. Application No.

PCT/US 99/15827

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/15 A61K31/4035 A61K31/496 A61K31/551 A61P9/10  
A61P7/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 519 035 A (MAIESE KENNETH ET AL) 21 May 1996 (1996-05-21)  claims 1,2  —  -/-	1-6, 9-11, 24, 25, 30, 31, 46-51, 54-56, 69, 70, 75, 76, 91-98, 167-176



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "a" document member of the same patent family

Date of the actual completion of the international search

1 February 2000

Date of mailing of the international search report

18/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Veronese, A

# INTERNATIONAL SEARCH REPORT

Intern:      Serial Application No

PCT/US 99/15827

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>EP 0 457 295 A (UNIV ROCKEFELLER) 21 November 1991 (1991-11-21)</p> <p>claims 1,6</p>	<p>1-6,21, 22,24, 25,30, 31, 46-51, 66,67, 69,70, 75,76</p>
X	<p>WO 95 14465 A (UNIV JOHNS HOPKINS ;US OF AMERICA NATIONAL INST O (US)) 1 June 1995 (1995-06-01)</p> <p>claims 1,8</p>	<p>1-6,21, 25,30, 46-51, 66,69,70</p>
P,X	<p>US 5 830 910 A (MATTSON MARK P) 3 November 1998 (1998-11-03)</p> <p>the whole document</p>	<p>1-6,21, 22,24, 25,30, 46-51, 66,67, 69,70,75</p>
X	<p>EP 0 654 266 A (ASAHI CHEMICAL IND) 24 May 1995 (1995-05-24)</p> <p>* See page 6, line 28: 1-(5-isoquinolinesulfonyl)homopiperazine (Fasudil) * claims</p>	<p>1-6,11, 16-18, 30, 46-51, 56, 61-63, 75, 131-137, 167-175</p>
X	<p>EP 0 187 371 A (ASAHI CHEMICAL IND ;HIDAKA HIROYOSHI (JP)) 16 July 1986 (1986-07-16)</p> <p>See page 8, line 9: 1-(5-isoquinolinesulfonyl)homopiperazine (Fasudil) * page 2, line 7-10</p> <p style="text-align: center;">-/-</p>	<p>1-6, 11-18, 30, 46-51, 56-63, 75, 131-137, 167-175</p>

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/15827

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online!  CHEMICAL ABSTRACTS SERVICE, COLUMBUS,  OHIO, US  HIDAKA, HIROYOSHI ET AL: "Preparation of  naphthalenesulfonylhomopiperazine and  -piperizine derivatives as cardiovascular  agents"  retrieved from STN  Database accession no. 107:154355 CA  XP002128508  * See compound RN= 109376-83-2 (ML-7) *  abstract  &amp; JP 62 087581 A (HOKURIKU PHARMACEUTICAL  CO., LTD., JAPAN)  22 April 1987 (1987-04-22)</p>	<p>1-6,  11-18,  30,31,  46-51,  56-63,  75,76,  91-98,  131-137,  167-176</p>
X	<p>PATENT ABSTRACTS OF JAPAN  vol. 012, no. 219 (C-506),  22 June 1988 (1988-06-22)  &amp; JP 63 017870 A (HOKURIKU SEIYAKU CO  LTD;OTHERS: 01),  25 January 1988 (1988-01-25)</p> <p>abstract</p>	<p>1-6,  11-18,  30,  46-51,  56-63,  75,  131-137,  167-175</p>
P,X	<p>WO 98 34615 A (VINER NORMAN M ;SYNAPSE  PHARMACEUTICALS INTERN (CA))  13 August 1998 (1998-08-13)</p> <p>claims 1,8,38</p>	<p>1-6,  11-18,  30,31,  46-51,  56-63,  75,76,  91-98,  131-137,  167-176</p>
A	<p>WO 91 06293 A (UNIV VERMONT)  16 May 1991 (1991-05-16)</p> <p>the whole document</p> <p style="text-align: center;">-/-</p>	<p>1-6,  11-18,  30,31,  46-51,  56-63,  75,76,  91-98,  131-137,  167-175</p>

**PCT/US 99/15827**

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/15827

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LAUFS U ET AL: "Upregulation of endothelial nitric oxide synthase by HMG CoA reductase inhibitors." CIRCULATION, (1998 MAR 31) 97 (12) 1129-35., XP000870148 the whole document	1-216
X	FUJITA, SATOSHI ET AL: "Effects of vasodilators and perfusion pressure on coronary flow and simultaneous release of nitric oxide from guinea pig isolated hearts" CARDIOVASC. RES. (1998), 38(3), 655-667 , XP000870320	1-6, 16-18, 24,25, 30,31, 46-51, 61-63, 69,70, 75,76, 131-137, 167, 173-176 1-216
Y	* See butanedione monoxime, results, discussion * page 656, column 2, <u>last paragraph</u>	



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/15827

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim(s) 1-216  
is(are) directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☒ Claims Nos.: 1-216 (partially)  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99 15827

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-216 (partially)

The present application has no less than 216 claims.

This presentation makes it difficult, if not impossible, to determine the matter for which protection is sought, to the extent that a complete search is not possible.

The number of claims has to be reasonable in consideration of the nature of the invention claimed. Obviously no hard and fast definition can be accorded to the term reasonable, but this does not mean that the requirement is wholly devoid of meaning and legal effect.

What is or what is not reasonable depends on the facts and circumstances of each particular case. In deciding the matter, regard has to be had to the interest of the relevant public, since patents should not be allowed to erect a legal maze or smoke screen in front of potential users of the inventions to which they lay claim. Patent claims, taken singly as well as in totality, must be clear and concise (Art. 6 and Rule 6.1 (a) PCT), in order to enable such potential users to ascertain, without undue burden, let alone recourse to litigation, whether their planned commercial use is likely to infringe the patent monopoly. Having to construe the 216 claims and to form a valid and commercially useful opinion on whether or not any one of them could prevent or hinder the commercial activities, must in the nature of things impose a severe and totally undue burden on the public. It follows that the present application as it stands falls foul of the clear provisions not only of Rule 6.1(a) PCT but also of Article 6 PCT.

The present application claims the use of an agent that disrupts actin cytoskeletal organization to increase endothelial cell Nitric Oxide Synthase activity, with beneficial effects in the treatment of a large number of diseases (cardio-circulatory, neurodegenerative, renal diseases and others). A direct relation between the mechanism of action which has been discovered, and the very broad spectrum of diseases mentioned in the application, is however not demonstrated (Art. 6 PCT). It is clear that a complete search is virtually impossible in this situation, because the compounds are defined by means of a pharmacological /enzymatic mechanism instead of by means of structural features.

The use of some of above mentioned compounds falling under that mechanism is already very well known in relation to the treatment of the same diseases mentioned by the inventors; this is also indicated by the presence of a number of disclaimers in the claimed subject matter. The literature in this regard is so vast that a complete retrieval of the pertinent documents is not possible for practical reasons.

Additionally, the use of a compound for the manufacture of a medicament for the treatment of a specified disease, once disclosed, cannot be patented again under the guise of another or newly specified pharmacological mechanism. In fact, the discovery of such a new way of action is not an invention under Rule 39.1(i) PCT, as the technical effect obtained remains the same (treating the same disease). In the present case the discovery of the mechanism of action, or an alternative mechanism of action does not add a new or better technical effect to the known treatment. The technical effect being identical the use is not

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99 15827

### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

changed by the discovery of the (alternative) mechanism. Moreover, the experimental data which have been presented in the description deal mostly with compounds which are actually disclosed (simvastatin, lovastatin).

Only Example 27 (page 57) deals with the pertinent compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, therefore, for only a very small proportion of the compounds/methods claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds which are exemplified in Example 27, with due regard to the idea underlying the application, and in view of the vast prior art, with particular regard to patent literature.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/15827

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5519035	A	21-05-1996	NONE	
EP 0457295	A	21-11-1991	AT 151287 T	15-04-1997
			CA 2042668 A	17-11-1991
			DE 69125523 D	15-05-1997
			DE 69125523 T	14-08-1997
			DK 457295 T	28-04-1997
			ES 2102986 T	16-08-1997
			GR 3024032 T	31-10-1997
			JP 7025786 A	27-01-1995
			US 5385915 A	31-01-1995
			US 5538983 A	23-07-1996
WO 9514465	A	01-06-1995	NONE	
US 5830910	A	03-11-1998	NONE	
EP 0654266	A	24-05-1995	JP 2893029 B	17-05-1999
			JP 6056668 A	01-03-1994
			US 5747507 A	05-05-1998
			CA 2141969 A	17-02-1994
			WO 9403171 A	17-02-1994
			KR 148645 B	02-11-1998
EP 0187371	A	16-07-1986	JP 1795364 C	28-10-1993
			JP 5003851 B	18-01-1993
			JP 61227581 A	09-10-1986
			JP 1789975 C	29-09-1993
			JP 4081986 B	25-12-1992
			JP 61152658 A	11-07-1986
			AT 64598 T	15-07-1991
			US 4678783 A	07-07-1987
JP 62087581	A	22-04-1987	JP 1849895 C	21-06-1994
			JP 5058433 B	26-08-1993
JP 63017870	A	25-01-1988	JP 2528451 B	28-08-1996
WO 9834615	A	13-08-1998	US 5900418 A	04-05-1999
			US 5981549 A	09-11-1999
			US 5902816 A	11-05-1999
			US 5824684 A	20-10-1998
			US 5760049 A	02-06-1998
			US 5916903 A	29-06-1999
			AU 5977598 A	26-08-1998
			ZA 9800960 A	17-08-1998
WO 9106293	A	16-05-1991	US 5162374 A	10-11-1992
WO 9741857	A	13-11-1997	US 5856360 A	05-01-1999
			AU 2740097 A	26-11-1997
			CA 2251969 A	13-11-1997
			CZ 9803444 A	14-04-1999
			EP 0904078 A	31-03-1999
			NO 985116 A	02-11-1998
WO 9947153	A	23-09-1999	AU 3107599 A	11-10-1999

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/15827

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9900117 A	07-01-1999	AU 8173098 A	19-01-1999